

S2ESWCTS  
DUSTY MAC

# **Barium-Strontium relationships**

**possible geochemical tool in search for ore bodies**

**H V Warren, B N Church, K E Northcote**


Reprinted from  
**Western Miner**  
issue of April, 1974

The study of the relative and absolute concentration of barium and strontium in country rocks may prove useful in searching for mineral deposit extensions. Data are given from two properties (silver and copper) in British Columbia.

## Barium-Strontium relationships

### possible geochemical tool in search for ore bodies

H V Warren, B N Church, K E Northcote



H V Warren



B N Church



K E Northcote

**Dr Harry V Warren** is a widely-recognized authority on geochemistry who has contributed many technical articles to *Western Miner*. He continues to be very active on the mining scene, though he retired in 1973 from the post of professor of geology at the University of British Columbia (UBC) which he held for many years.

**Dr B Neil Church** has worked with the Geological Survey of Canada, the governments of Saskatchewan and Manitoba, and with private industry. For several years he has been regional geologist with the BC Department of Mines and Petroleum Resources, specializing in mineralization related to volcanic deposits.

**Dr K E Northcote** has had wide geological experience in Canada and Colombia (S America), and has been for several years with the BC Department of Mines and Petroleum Resources; he has been engaged in geological mapping on Vancouver Island.

Much is known about the distribution and concentration of barium and strontium in minerals and rocks but relatively little use has been made of the wide variations that are found in eruptive rocks, particularly in the vicinity of some ore bodies.

This paper indicates how the study of the relative and absolute concentrations of barium and strontium in country rocks may prove useful in searching for mineral deposit extensions such as at two well known British Columbian properties, the *Dusty Mac*, a silver occurrence in the southern Okanagan, and *Island Copper* mine in northern Vancouver Island.

Rock geochemistry has recently become important as a standard exploration tool and is especially useful in heavily drift-covered areas.

Initially, rock geochemistry was concerned with the concentrations in apparently barren rock of the metals which were being sought, such as copper, zinc, molybdenum, and gold. More recently attention has been turned to other elements whose distribution in rocks near mineral deposits may be useful because they provide haloes more easily recognizable than those provided by the metals which are sought. Barium and strontium, the elements with which this paper is primarily concerned, are both relatively abundant and readily determined in most rocks.

#### Concentrations of Ba and Sr in eruptive rocks

The concentrations of barium and strontium in various rocks and minerals have been documented by many writers, and there is a general agreement between most of them, including Goldschmidt (1954), Rankama and Sahama (1949), Turekian and Wedepol (1961), Holmes and Harwood (1937), and Hawkes and Webb (1962).

Strontium is present in both plagioclase and in alkali feldspar, but significant amounts of barium are found only in alkali feldspar and biotite.

One of us (BNC) has studied the concentrations of barium and strontium in some feldspars and feric minerals. Biotite and anorthoclase contain the greatest amounts of barium but the plagioclases tend to show more strontium. Unfortunately only seventeen determinations were made and consequently more detailed work might modify these findings although they do appear to be in accordance with the results obtained by other workers.

Goldschmidt suggested an average concentration of 430 ppm (parts per million) of barium for the igneous rocks of the earth's crust, and Hawkes and Webb gave 640. All these writers have stated that in the sequence from gabbros, basalts, and anorthosites to granites and syenites the barium concentrations rise from something under 100 ppm to well in excess of 1000 ppm, being especially high in some syenites, trachytes, and phonolites.

In basic rocks strontium generally exceeds the barium content but the reverse is normally found in granites, syenites, and related volcanic rocks. Grove (1971) drew attention to barren vein material relatively enriched in barium in a mineralized zone extending about 60 feet above known bonanza ore shoots in the *Silbak Premier* vein system, and suggested rock geochemistry would be a useful exploration method in that area.

Grove (1971) drew attention to barren vein material relatively enriched in barium in a mineralized zone extending about 60 feet above known bonanza ore shoots in the *Silbak Premier* vein system, and suggested rock geochemistry would be a useful exploration method in that area.

Grove (1971) drew attention to barren vein material relatively enriched in barium in a mineralized zone extending about 60 feet above known bonanza ore shoots in the *Silbak Premier* vein system, and suggested rock geochemistry would be a useful exploration method in that area.

#### I. *Dusty Mac*

The *Dusty Mac* property is situated one mile east of the south end of Skaha Lake and the village of Okanagan Falls in the southern part of the Okanagan Valley. The property has been described in detail (Church, 1973) and only a brief resume of the general geology is given here (Fig 1).

**General geology**

In the vicinity of the Dusty Mac silver-gold mineralization three rock assemblages are important. They are parts of the Tertiary Marama and White Lake Formations, and a tectonic breccia characterized by quartz veining, much silicification and gossan.

The zone of potential economic mineralization is where silicification is greatest on what is believed to be the limb of a faulted syncline in rock identified as part of the White Lake Formation (Fig 2 & Fig 3).

Silicification and gossan are the best local guides in looking for ore but outcrop is sparse in much of the area of best potential. Fortunately many rock samples taken both from the available outcrops and from drill cores have been available for study and trace element analysis.

Barium, strontium, chromium, nickel, silver, and lead were determined at the Department of Geological Sciences, University of British Columbia, by Don Marshall using a large Hilger spectrograph. Tellurium, arsenic and quartz analyses were carried out in the laboratories of the British Columbia Department of Mines and Petroleum Resources, in Victoria.

Five tables of data were prepared from these results: A: Seventeen representative samples of the Marama Formation. B: Eleven from the White Lake Formation. C: Nine from the 'ore', obtained by drilling. D: Six from within fifty feet of the 'ore' as outlined by drilling. E: Seventeen samples from cores taken from drill holes unsuccessful in finding ore. The locations from which all these samples were taken are given in maps (Fig 3 & 4).

**Discussion of Dusty Mac results**

On the basis of the data contained in table I some conclusions may be drawn:

(1) In only one sample (DDH 14-8') with a quartz content of more than 50% is the silver content below 15 ppm, or 0.5 oz per ton.

(2) In only one sample (DDH 8-19') with a quartz content of less than 50% is the silver content more than 25 ppm, or 0.83 oz per ton.

(3) In general the 'no ore' holes are all in non-silicified White Lake Formation.

(4) The low barium and strontium values in 'ore' cannot readily be explained simply by the abundance of quartz and a relative deficiency of alkali feldspar. Only in the 'ore' samples are strontium values greater than barium. Only in basic and ultra mafic rocks could this be considered normal.

(5) Silver values cannot be related directly to either lead or arsenic and it seems probable that some lead and silver can be attributed to telluride minerals.





Fig. 1. Dusty Mac: general location map

Fig. 2. Dusty Mac: faulted synclinal structure



## II. Island Copper and Red Dog

Both properties are situated in the Rupert Inlet — Cape Scott Map-Area in the northern part of the Nanaimo Mining Division of British Columbia.

Two eruptive rock units, the Karmutsen Formation and the Bonanza Subgroup, are important in this discussion. Northcote (1970) has described the general geology of the area and the following two paragraphs are summarized from his report.

The Karmutsen Formation is of Triassic age and the rocks are characteristically medium to dark green-grey, interbedded massive porphyritic and amygdaloidal flows of basaltic composition. Tuff is interbedded with flows and some pillow breccia, and poorly developed pillows occur.

The Bonanza Subgroup ranges from Upper Triassic to Upper Lower Jurassic. We are concerned with an upper volcanic unit, made up of bedded and massive tuff, lapilli tuff, tuff breccia, and some interbedded lava flows. These volcanic rocks range from basaltic andesite to rhyolite in composition.

Northcote selected the following samples as representatives of these rock units: A, 7 Karmutsen; B, 12 Bonanza (fresh); C, 6 Bonanza (altered); D, 3 Intrusive rocks; E, 2 Bonanza (Island Copper ore zone); F, 1 Bonanza (fringe of Island Copper ore); G, 5 Bonanza (Red Dog mineralization). The locations of these samples are shown on Figure 5.

Mr John Lamb, Senior Geologist at Island Copper Mine, kindly sent seven selected samples representing the ore and material on the fringe, respectively, H and I in Table 2.

All of the above samples were analyzed for barium, strontium, chromium, nickel, lead, silver, and copper by Don Marshall at the University of British Columbia.

These results have been tabulated in Table 2.

### Discussion of Island Copper and Red Dog results

It can be seen from Table 2E, 2G, and 2H that the barium concentrations fall below those of strontium in the samples taken from ore zones in marked contrast to fringe samples 2F and 2I. Table 2B, for unaltered Bonanza rocks shows average barium values are greater than strontium values.

Table 2C, for altered Bonanza rocks, shows two groups of contrasting barium-strontium ratios of six analyses listed, four have barium values less than strontium. The remaining two samples are enriched in barium relative to strontium. It is of interest to note that three of the four samples with low barium-strontium ratios can be related to alteration associated with copper mineralization but were not taken from ore zones.

The two samples with high barium-strontium ratios show epidotization and/or pyritization but are not known to be associated with copper mineralization.

Barium is consistently lower than strontium in all Karmutsen samples, presumably because of the more basic nature of these rocks.

From the data presented here it appears that each eruptive rock unit exhibits a characteristic barium-strontium relationship. This relationship, on occasion is varied, in places greatly, where there is mineralization nearby. Further, it is suggested that rocks which have undergone different types of alteration will

exhibit different barium-strontium ratios.

In the examples given in this paper, the barium-strontium relationships in the ore bodies are reversed from those of the host rock units.

Although this paper has considered only a few examples, and no final conclusions can be made, enough evidence has been accumulated to justify further studies.

Barium and strontium are two elements which lend themselves admirably to bio-, pedo-, equally well as to litho-geochemistry. This fact may be important in areas where extensive overburden makes rock geochemistry impractical.




Fig. 3. Dusty Mac: location of samples (Table 1)

**Acknowledgements**

One of us (HVW) is indebted to The National Research Council (Grant No.1805), a Canada Council Grant (Kilam Award), and The Geological Survey of Canada for financial support for investigations which included this project. The study was greatly facilitated through use of the rock library system of the British Columbia Department of Mines and Petroleum Resources.

**REFERENCES**

Grove, E W (1971). 'Geology and mineral deposits of the Stewart Area'. Bull. No.58. BCDM, 219p.  
 Goldschmidt, V M (1954). 'Geochemistry'. The Clarendon Press Oxford, pp243-256.  
 Rankama, K and Sahama Th J (1949). 'Geochemistry'. University of Chicago Press, Chicago, pp470-477.  
 Hawkes, H E, and Webb, J E (1962). 'Geochemistry in mineral exploration'. Harper and Row, New York, pp360-361.  
 Holmes, A and Harwood H F (1937). 'The petrology of the volcanic area of Bufumbira'. Geol. Sur. of Uganda Memoir No.111, Part II pp48-49.  
 Church, B N (1973). Geology of the White Lake Basin; BCDM Bull. 61, 120p.




Fig 4. Dusty Mac: location of samples (Table 1)

Turekian, K K, and Wedepol, K H (1962). 'Distribution of the elements in some major units of the earth's crust. pp175-191.  
 Northcote, K E (1970). 'Rupert Inlet-Cape Scott map area'. Geology, exploration, and mining, BCDM pp254-269.






Table 1. Some trace elements and quartz in various samples (chemical data in ppm)

A. Marama Formation (Rhyolite - Dacite)									C. Mixed Quartz Breccia (Silver 25 ppm or more)									
Sample No.	Ba	Sr	Cr	Ni	Pb	Ag	As	Quartz %	Sample — DDH	Ba	Sr	Cr	Ni	Pb	Ag	Te	As	Quartz %
57	100	400	15	5	7	—	4	46	9-21'	250	175	3	<4	50	25	<15	5	93
59	300	200	8	<4	5	—	8	35	9-26'	400	250	30	<4	30	500	<15	7	100
64	500	300	4	<4	10	—	3	40	8-19'	500	500	2	<4	500	50	23	6	35
66	500	300	2	<4	10	—	1	42	8-26'	350	400	5	—	100	100	<15	4	100
101	150	400	10	4	8	—	4	31	10-24'	<100	300	3	30	400	500	375	148	75
102	900	300	10	4	10	—	8	35	10-31'	<100	500	<1	15	150	200	<15	8	100
105	600	350	8	—	8	—	2	49	11-23'	<100	100	<1	—	35	100	<15	5	100
108	500	300	9	<4	9	—	3	30	12-32'	500	500	4	—	50	100	<15	6	90
110	400	400	10	7	8	—	1	42	13-25'	300	200	<1	—	800	1000	105	10	66
111	400	300	8	4	7	—	2	29	Sum	<2500	2925	<50	<57	2115	2527	<593	197	759
131	200	700	10	4	8	—	5	30	Average	<280	325	<6	<7	235	285	<65	22	84%
137	600	500	5	<4	7	—	5	31	D. Within 50 feet of 'ore' zone (silver less than 25 ppm)									
146	400	300	2	—	10	—	2	47	Sample — DDH	Ba	Sr	Cr	Ni	Pb	Ag	As	Quartz %	
148	400	300	2	<4	8	—	1	40	9-47'	1000	400	4	—	8	—	6	41	
154	900	500	10	4	10	—	2	31	8-51'	1000	400	1	<4	10	—	5	29	
155A	300	300	10	5	10	—	2	40	11-87'	1000	400	<1	4	10	—	10	38	
158	400	200	10	5	8	—	1	48	12-73'	2000	400	1	<1	15	—	12	32	
Sum	7550	6050	133	<66	143	—	54	646	13-7'	2500	500	20	—	10	—	7	47	
Average	445	355	8	<4	8	—	3	38%	13-69'	1000	500	2	—	15	—	10	37	
B. White Lake Formation (Andesite)									Sum	8500	2600	<29	<4	68	—	50	224	
Sample No.	Ba	Sr	Cr	Ni	Pb	Ag	As	Quartz %	Average	1420	415	<5	<1	11	—	8	37%	
56	1800	500	250	40	15	—	8	20	E. Unsuccessful drill hole samples									
72	1000	500	50	8	10	—	3	19	Sample — DDH	Ba	Sr	Cr	Ni	Pb	Ag	As	Quartz %	
82	2000	1000	100	15	30	—	11	20	14-8'	1500	700	12	—	12	10	11	83	
87	2000	600	125	25	15	—	8	26	14-101'	3000	700	2	—	8	10	11	28	
96A	1000	1000	150	30	20	—	3	24	15-14'	2000	1000	100	30	20	—	8	24	
96B	1350	950	100	25	12	—	5	24	15-96'	2000	600	2	—	18	—	11	29	
147	6000	700	70	12	10	—	2	31	18-31'	1000	500	1	—	15	—	13	39	
162	1000	400	100	20	10	—	1	20	23	2000	1000	40	4	15	—	13	14	
165	2000	650	100	25	15	—	1	25	24-52'	2000	800	100	20	15	—	—	—	
168	1000	500	100	20	10	—	1	20	39-75'	1000	400	<1	—	8	2	16	—	
176	2000	700	70	10	18	—	2	21	39-145'	1000	500	3	—	10	—	13	27	
Sum	21150	7500	1215	230	165	—	45	250	40-55'	1500	1000	2	—	18	—	11	18	
Average	1920	680	110	21	15	—	4	23%	40-140'	1750	700	2	—	20	—	12	24	
— not detectable by analytical method employed									42-18'	4000	1000	200	30	15	—	15	17	
<b>SUMMARY</b>									42-105'	1500	500	30	—	10	—	13	23	
	Ba	Sr	Cr	Ni	Pb	Ag	As	Quartz %	45-31'	3500	1500	30	4	18	—	6	25	
A. Marama	445	355	8	<4	8	—	3	38	45-45'	2500	1000	35	20	20	—	5	15	
B. White Lake	1920	680	110	21	15	—	4	23	45-73'	2500	2000	125	20	15	—	10	13	
C. 'Ore'	<280	325	<6	<7	235	285	22	84	45-155'	2000	1000	15	<4	40	—	7	20	
D. Near 'ore'	1420	415	<5	<1	11	—	8	37	Sum	34750	14900	700	132	277	22	175	399	
E. No 'ore' holes	2040	875	40	8	15	—	10	25	Average	2040	875	40	8	15	1	10	25%	

# PRELIMINARY GEOLOGICAL MAP RUPERT INLET - CAPE SCOTT AREA

GEOLOGY BY K.E. NORTHCOTE

## LEGEND

-  **INTRUSIVE ROCKS**  
VARIED COMPOSITION FROM DIORITE TO GRANITE AND INCLUDES PORPHYRITIC PHASES
-  **LOWER CRETACEOUS SEDIMENTARY ROCKS**  
CONGLOMERATE, SANDSTONE, SILTSTONE, SHALE, CARBONACEOUS HORIZONS.
-  **BONANZA SUBGROUP**  
UPPER VOLCANIC UNIT; LARGELY PYROCLASTIC TUFF, LAPILLI TUFF AND TUFF BRECCIA OF ANDESITE AND BASALT COMPOSITION WITH SOME BASALT AND RHYODACITE FLOWS AT THE TOP OF THE UNIT.  
LOWER SEDIMENTARY UNIT; THIN BEDDED ARGILLACEOUS AND CARBONACEOUS LIMESTONE, CALCAREOUS SHALE AND SILTSTONE AND GREYWACKE.
-  **QUATSINO FORMATION**  
LIMESTONE, MEDIUM TO THICK BEDDED
-  **KARMUTSEN FORMATION**  
BASALTIC AMYGDALOIDAL AND MASSIVE FLOWS, INTERBEDDED TUFF, SOME PILLOW BRECCIA AND POORLY DEVELOPED PILLOWS. THIN LIMESTONE BEDS NEAR TOP OF FORMATION

## SYMBOLS







- CONTACTS:
- KNOWN 
  - APPROXIMATE 
  - ASSUMED 
  - LINEAMENTS FROM AIR PHOTOGRAPHS  
SOME OF THESE ARE KNOWN TO REPRESENT FAULTS 
  - BEDDING 
  - MINERAL DEPOSITS 

Fig 5. Location of representative rock samples (Table 2)





Table 2. Some trace elements in various sam 1 ppm)

A. Karmutsen								D. Intrusive rocks							
Sample no.	Ba	Sr	Cr	Ni	Pb	Ag	Cu	Sample No.	Ba	Sr	Cr	Ni	Pb	Ag	Cu
PN 70 131 A	100	500	40	10	3	2	40	KN 70-2-1	200	700	200	20	15	2	500
PN 70 153	100	200	200	100	20	2	1000	PN 70-167B	200	500	150	30	100	2	500
PN 70 140	100	200	200	100	2	2	100	CN 70-127A	100	100	400	150	80	2	800
KN 68 244	100	500	300	50	25	2	1000	CN 70-132	100	80	150	18	10	2	300
KN 68 295	100	500	400	100	7	2	500	Sum	600	1380	900	218	205	8	2100
CN 70 83	100	400	1000	150	5	2	1000	Average	150	345	225	55	51	2	525
KN 68 70	200	800	200	40	10	2	500	E. Bonanza (Island Copper ore zone)							
Sum	800	3100	2340	550	72	14	4540	KN 68-93	100	400	200	30	8	2	500*
Average	110	445	335	80	10	2	650	KN 68-89	100	1000	100	30	10	2	200
B. Bonanza (fresh)								KN 68-78	100	1000	200	50	70	2	800
Sample no.	Ba	Sr	Cr	Ni	Pb	Ag	Cu	Sum	300	2400	500	110	88	6	1500
KN 68-41	400	400	200	50	5	2	400	Average	100	800	170	37	29	2	500
KN 68-22	2000	1000	500	50	5	2	500	* Pyrophyllite-dumortierite							
KN 68-18	200	800	300	100	18	2	1000	F. Bonanza (fringe of Island Copper ore)							
KN 68-66	1500	1000	200	40	35	2	600	KN 68-335	1500	400	200	15	4	2	200
KN 69-75	700	600	190	30	7	2	250	G. Bonanza (Red Dog mineralization)							
KN 69-74-I	400	400	200	60	2	2	150	KN 70-5	100	600	800	200	10	2	5000
KN 69-96-V	500	300	100	20	10	2	400	KN 70-6	100	500	1000	500	5	2	5000
KN 69-147	1000	800	100	30	15	2	400	KN 70-6-III	100	800	400	100	5	2	5000
KN 69-272A	3000	1000	50	10	10	2	200	KN 70-7-VIII	100	300	200	2000	60	2	5000
PN 70-178	200	300	180	25	100	2	300	Sum	400	2200	2400	2800	80	8	20000
CN 70-34	500	500	200	30	15	2	400	Average	100	550	600	700	20	2	5000
CN 70-69A	200	500	200	40	15	2	500	H. Bonanza (Island Copper ore) (Lamb)							
Sum	10600	7600	2420	485	237	24	5100	V 51	150	500	100	8	15	4	3000
Average	885	635	200	40	20	2	425	W 49	150	400	100	12	10	2	4000
C. Bonanza (altered)*								W 52	200	600	130	12	10	2	4000
KN 68-21	100	800	200	20	15	2	5000	Sum	500	1500	330	32	35	8	11000
KN 69-143	3000	600	180	20	20	2	500	Average	170	500	110	11	12	3	3670
KN 69-166A	5000	800	100	10	20	2	200	I. Bonanza (fringe of Island Copper ore) (Lamb)							
KN 69-202-I	100	300	100	15	15	2	200	Sample no.	Ba	Sr	Cr	Ni	Pb	Ag	Cu
CN 70-36-I	100	300	150	15	15	2	400	P 24	100	100	75	6	5	2	200
CN 70-162	100	400	200	40	8	2	5000	Q 23	300	100	80	10	5	2	1000
Sum	8400	3200	930	120	93	12	11300	P 22	400	150	110	10	15	2	1000
Average	1400	535	155	20	16	2	1883	Q 21	400	100	50	10	5	2	3000
Omitting								Sum	1200	450	315	36	30	8	5200
KN 69-143	400	1800						Average	300	110	80	9	7	2	1300
166A	100	450													

\* Epidotized, sericitized, pyrophyllitized, silicified and/or pyritized.