002504

082F NW 240

CONDOR, (IRON MASK, BOCO II)

82F/11E 8W 49 55 48 117 15 36

A.R. 16,247 : Orientation soil geoc & geol surveys Iron Mask, Condor, Boco I mineral claims Slocan Mining Division, Goldsmith L, 1987

- massive ZnS @ Ag values approx 5-10 oz/ton occur in a NE trending lode hosted in competent qte & argillite. width of lode structure .5m to fracturing across perhaps 50 m. "The latter dimension is more representative of the structure, while the .5m width where exposed in a pit is prob only 1 mineralized strand within the lode zone."
- soil geoc not effective in tracing strike extensions
- UCA Roi / Hewitt mines are 1.5 - 2.5 km to SW.
- Slocan Gp Charred & clastic clastic sediments : banded argillites qte & thin bedded limestone @ NW bedding trends & steep E dips. Folding generally oriented NW.
- A NW trending N dipping lode system on the Iron Mask approx 5-10 oz/ton Ag & 20% Zn. Lode structure @ strands of ZnS & gh are exposed in discontinuous outcrops along road cut. Fracturing assoc @ lode appears to occur across 50 m true width, ↓ in intensity to W.

- Attitude of (zone) is similar to that mirrored in VAN ROI & Haith