



- | | | |
|--|--|--|
| | | Carbonate rock - magnesite, dolomite etc. - replacement, often breccia |
| | | Feldspar and quartz-feldspar porphyry - dykes |
| | | Brown weathering basalt (andesite) - flows dykes sills |
| | | Cherry Creek intrusions - fine grained porphyritic diorite |
| | | Sugarloaf porphyritic diorite, usually contains hornblende |
| | | Ultra-mafic intrusions - perite, peridotite |
| | | Buff colored sediments - sandstone, argillite |
| | | Kamloops Group |
| | | Coarse grained porphyritic flows, dolerite |
| | | Fine grained amygl flows and poorly lithified tuff |
| | | Agglomerate and coarse well lithified tuff |
| | | Coldwater? Conglomerate |
| | | Iron Mask Batholith - diorite, minor pyroxenite and gabbro |
| | | Nicola Group Volcanics - Greenstone, agglomerate, minor tuff |
| | | Sediments - Argillite, lime argillite |
| | | Cache Creek Group - Argillite, limestone, chert |
| | | Micro-granite |


Miocene and/or younger

Miocene and/or Oligocene

Jurassic?

Upper Triassic

Permian & older



9219w 610c

NEW JERSEY ZINC EXPLORATION CO. (CANADA) LTD.

(Unpublished) **GEOLOGY**
Kamloops Area

Map accompanying report by E. Livingston

Scale 1" = 1/2 mi. Plot by J. M. Carr

DATE: Apr. 60
BY: E.L.

009650