CANADIAN SUPERIOR EXPLORATION LIMITED

DIAMOND DRILL RECORD

010405

HOLE NO: 72-2

LOCATION: SITKOM 326+00-E

AZIMUTH: 316°

DIP: -60°10'

LENGTH: 600 FT.

ELEVATION: 4700 ASL ± 50'

PROPERTY: CHATTAWAY EXPLORATIONS LTD

CLAIM NO: LEN 12

STARTED: April 3rd 1978

CORE SIZE: BQ

DATE LOGGED: April 6th, 1978. April 7th 1978

SECTION: —

COMPLETED: April 6th 1978

DIP TESTS: -45½° @ 250' -41½ @ 300'

LOGGED BY: JAY D. MURPHY

PURPOSE: TEST IP ANOMALY

FOOTAGE from	FOOTAGE to	DESCRIPTION	SAMPLE NO.	FOOTAGE from	FOOTAGE to	LENGTH
0.0	17.0	Casing. Shot in 8.6°				
17.0	176.1	Granodiorite: Chattaway variety, generally fresh, pinkish grey, med. grained, uniform, massive, well formed subhedral to euhedral mafics (hornblende & biotite as outstanding idiomorphic, hbde. slightly chloritized, otherwise little evidence of alteration or movement), minor pink K-feldspar.				
		65.1 - ½" gouge & fractured slip @ 25° w. parallel carb. str., previous 36" core "blocky"				
		76.2 - 24" dyke, top 2" splits adjacent to old shear @ 55°, balance of section med. grey, fine grained w. coarse mafic phenos. Fairly high percentage of mafics mainly as fine platey xl's in matrix, classified tentatively as Klitchen Brook "B".				
		1090-1295 - Shattered, core strongly broken up, blocky, abundant development of plastic gouge; up to 12", rock has more ground feel due to alteration of mafics				

SUPERIOR EXPLORATION LIMITED

DIAMOND DRILL RECORD

HOLE NO: 72-2

PAGE NO: 2

FOOTAGE from	FOOTAGE to	DESCRIPTION	SAMPLE NO:	FOOTAGE from	FOOTAGE to	LENGTH
17.0	171.7	Conglomerate: Chetaway Variety (contd)				
		129.0 – 135.5 (contd) sand grained alti of feldspar considered as recrystallization, attitude of core not definite but 2" gouge at start of section faulty detrite @ 50°, minor introduced carb. also @ 30-35°, final 2½" section consists of 1½" clayey gouge followed by 2" crushed section consisting of coarse rock fragments in matrix of rock flour, grades back to faulty fine to coarse granodiorite, showing nice end of section at about 50°,				
171.7	225.4	Granodiorite: Watches Brook Variety B relatively fresh, medium grained grey, slightly porphyritic w. med to coarse blebs in f.g. matrix, faintly massive homogeneous appearance, top def. w. Chetaway sharp @ 60° with 1" pk. feldspathic band, rock becomes progressively deformed due to recrystallization of feldspars as shatter zone is approached,				
		185.4 - 221.8 Shatter Zone; rock strongly fractured, brecciated and altered to gouge, shatter zone showing less evidence of movement than has been strongly recrystallized with resultant dark green color, start of section indistinct measured @ 40°, core showing dips at several angles to core axes, minor introduced carbonate also appreciable qtz as mineralized				

CANADIAN SUPERIOR EXPLORATION LIMITED

DIAMOND DRILL RECORD

HOLE NO: 72-2

PAGE NO: 3

FOOTAGE from	FOOTAGE to	DESCRIPTION	SAMPLE NO:	FOOTAGE from	FOOTAGE to	LENGTH	Cu.	MoS_2	Au.
171.7	223.4	Granodiorite: Klotaw Brook Variety B (contd							
		180.4-224.8 - Shatter Zone (contd) 'crackled' veins & fragments with gneiss; minor graphite along slip & mixed with clayey 'gouge' with numerous well defined slips between 25° 50° to core, predominant attitude 35-40° Minor Bornite & Molybdenite in qtz.							
		Sample: dark green soft mainly gouge & fragments	229	200.5	202.5	2.0	0.09	0.025	
		Sample: grey crackled qtz. w. Bornite & MoS_2 along narrow fractures, graphitic 'gouge' over 6" at start of section	230	202.5	204.5	2.0	0.47	0.345	
		Sample: mainly gouge w. qtz. frags.	231	204.5	206.5	2.0	0.38	0.036	
		Sample: 12" gouge followed by 'blocky' fractured section	232	206.5	211.5	5.0	0.11	0.172	
		Sample: 3' qtz. at 40° w. fair MoS_2 as disseminated flakes & coatings along contacts	233	211.5	213.0	1.5	0.07	0.158	
		Sample: strongly fractured, barren greenstone 7' ground core.	234	213.0	216.0	3.0	0.03	0.028	
		strongly altd (serecitized) section of shatter zone has little or no effect on magnetic in contrast to less altd sections.							
223.4	500.0	Granodiorite: Chatoway Variety, generally fresh appearance, med. grey color w. pinkish cast probably due to Kspar alter, normally uniform, massive w. faulted, fractured sections as noted, occasional reddish orange unidentified min. as v. fine stks. ($Zn(O_2^2)$)							
		228.0-233.0 - 74" lost core							
		233.0 - 6" strongly graphitic 'gouge' in fault zone @ 45°							

CANADIAN SUPERIOR EXPLORATION LIMITED

DIAMOND DRILL RECORD

HOLE NO: 72-2

PAGE NO: 4

FOOTAGE from	FOOTAGE to	DESCRIPTION	SAMPLE NO:	FOOTAGE from	FOOTAGE to	LENGTH
225.4	500.0	GRANODIORITE Chatonby Variety (Cont'd)				
		236.0 - Fault zone consisting of 24" rock flour fragments & gouge followed by 16" strongly graphitic gouge, apparent attitude 30°				
		2325 - 30" section strongly altd (serpentised ?) to dark green color, soft, chloritic, carbonaceous, grades to normal granodiorite				
		2620 - 268.0: 3.8' lost core due to reported 'round section'				
		285.5 - 6" gouge developed				
		285.5 - 330.5 Shattered zone, very blocky with some development of fault gouge & slickensides, otherwise rock relatively unaltd, strong development of gouge over final 24", attitude of movement poorly defined, numerous minor slips @ 50°				
		460.5 - 481.0 - 'blocky' section with numerous irregular dry fractures @ 35-65° max indicative of tension rather than movement				
		464.5 - 24" fault zone strongly brecciated and clayey, movement appears 45-50° to core				
		499.5 - 18" fault zone, strongly brecciated & clayey, attitude not apparent,				
	500.0	End of Hole				