



**M. J. MINERALIZED AREA**  
**BEAR RIVER RIDGE - SKEENA M. D.**  
 1 inch = 100 Feet

- LEGEND**
- ICE and SNOW
  - GLACIAL MORAIN
  - QUARTZ-FELSITE / SSURE VEINS - Width & Sample Location
  - LAMPHOPHYRE DYKES
  - FELDSPAR PORPHYRY INTRUSIVES
  - NASS FORMATION - Banded Amphiboles & Quartzites
  - SALMON RIVER FORMATION - Conglomerates, Breccia
  - BEAR RIVER FORMATION - Volcanic Flows & Tuffs

		ASSAYS					
No.	Vein	Width	Au. oz.	Ag. oz.	Cu. %	Pb. %	Zn. %
327	1	18"	Trace	Trace	N/A	Trace	Trace
328	1	42"	Trace	0.10	0.05	Trace	Trace
329	1	25"	Trace	0.05	0.07	Trace	Trace
330	2	80"	0.018	1.30	0.32	2.65	Trace
331	2A	39"	0.01	0.50	0.07	1.92	Trace
332	2B	23"	0.005	0.60	0.06	2.60	N/A
333	3A	150"	Trace	0.10	N/A	N/A	N/A
334	3B	90"	0.005	Trace	N/A	0.10	Trace
335	6A	34"	0.005	0.50	0.12	Trace	Trace
336	6A	72"	0.005	0.25	0.12	Trace	Trace
337	6A	30"	0.03	9.30	0.10	17.05	11.75
338	6A	73"	0.01	2.20	0.20	4.70	5.45
339	6B	72"	0.01	0.25	0.10	0.40	Trace
340	7A	72"	0.01	0.45	0.15	1.15	Trace

Based on Triangulation and traverses by compass, tape and aneroid. To accompany report by W. N. Plumb, P. Eng.  
 November, 1956  
 W. N. Plumb