

D.H. No.	Intersection	Depth	8 Cu	8 Mo
PH - 1	50' to 400'	340'	0.21	0.007
PH - 2	40' to 220'	180'	0.19	0.005
PH - 3	10' to 200'	190'	0.12	0.005
PH - 4	30' to 300'	270'	0.10	0.007

With the limited amount of information available to date it appears that there is a north northwesterly trending zone of mineralized granodiorite that may average 0.28% Cu and 0.012% Mo. This is bounded on either side by material that may run in the 0.10 - 0.15% Cu and 0.005% Mo.

The quartz porphyry on the east side of the area is considered to be unfavourable.

An attempt was made to calculate reserves only including information from drill holes that have intersections that average greater than 0.20% Cu. Projections from drill holes are a maximum of 200 feet. A summary of these reserves are as follows:

D.H. NO.	Tons	<u> 8 Cu</u>	<u>% Mo</u>
81 - 1	1,300,000	0.25	.004
PH - 1	2,800,000	0.21	.007
A - 1	2,200,000	0.23	.007
81 - 2	11,100,000	0.29	.017
72 - 2	1,300,000	0.284	_
72 - 1	2,000,000	0.22	.005
	20,700,000	0.28	0.012

These blocks are shown on the accompanying plan and vertical sections.

It is interesting to note that the deepest hole (81 - 2) has the best grade material throughout with the best intersection from 588' to 888' averaging 0.393% Cu and 0.029% Mo.

- 2 -









