

da and BP Canada are strongly committed to environmental protection and have initiated base line research, to evaluate the environmental impact of mine development, as part of the Project Registration Process in Newfoundland.

The Duck Pond discovery represents a 10-year commitment by Noranda and BP Canada to further exploration and mine development. It illustrates the positive results of dedication and persistence to a favourable geological environment.

... Dan MacInnis
Colin MacKenzie
Gerry Squires

**CU-AU PORPHYRY WORKSHOP
April 5, 1989, Vancouver**

The following articles are the second installment of articles presented at the Cu-Au Porphyry workshop in Vancouver. Please refer to The Gangue No. 28 for the first three articles.

**Iron Mask Batholith
& Associated Fe-Cu-Au Deposits**

The Lower Jurassic Iron Mask Batholith is a composite alkaline intrusion located at the north end of the Nicola Volcanic Belt immediately west of Kamloops in south-central British Columbia. The batholith consists of two major bodies: the larger Iron Mask pluton to the southeast and the smaller Cherry Creek pluton to the northwest.

Extensive Cu-Au porphyry deposits of the alkalic class are hosted by later phases of the batholith, and include Afton (24.4 Mt @ 0.91% Cu and 0.020 oz/t Au), Ajax (27Mt @ 0.46% Cu and 0.01 oz/t Au), Big Onion (3.6Mt @ 0.71% Cu and 0.012 oz/t Au), Pothook (2.6Mt @ 0.35% Cu and 0.021 oz/t Au), and Crescent (1.3Mt @ 0.46% Cu and 0.006 oz/t Au).

Extensive zones of pervasive or vein controlled sodium silicate (albite), potassium silicate (potassium feldspar), propylitic (chlorite, epidote, carbonate and pyrite) and carbonate alteration (ankerite) are associated with these

deposits.

The Iron Mask Pluton consists of four major intrusive phases, the Iron Mask Hybrid, Pothook, Sugarloaf and Cherry Creek units, which range in composition from mafic agmatite and coarse grained gabbro to microsyenite porphyry. Field relationships, and major element geochemistry indicate that all four phases resulted from fractional crystallization of an original gabbroic magma under conditions of high oxidation and continuous crystallization of magnetite. K-Ar age dates of all four phases range from 194 to 2046 Ma, supporting these interpretations.

Emplacement of the Iron Mask and Cherry Creek plutons was controlled by a major northwest trending fault system and the plutons were emplaced at an epizonal to subvolcanic level in an active structural zone. The batholith is essentially coeval with, and is considered to be co-magmatic with, Nicola Group volcanic rocks.

Emplacement of the younger phases of the batholith, particularly the Sugarloaf unit and the more felsic and porphyritic phases of the Cherry Creek unit, was dominated by existing northwesterly trending structures, and these units are confined to the faulted edges of the Iron Mask pluton or along major cross-cutting faults.

Copper-gold mineralization is widespread throughout the batholith and closely associated with the Sugarloaf and Cherry Creek units. Bodies of Cherry Creek micromonzonite and microsyenite explosion breccia are particularly good hosts for mineralization. Field relationships indicate that copper-gold mineralization was deposited by structurally controlled hydrothermal systems that developed during the final stages of crystallization of Iron Mask magma. K-Ar age dates of hydrothermal biotite coincide with ages from primary biotite, supporting this interpretation. Magnetite lodes hosted by the Cherry Creek unit are titaniferous and apatite rich, and are considered to be immiscible segregations from the differentiating alkalic magma.

Note: Material presented in this paper is based on studies by the author, K.E. Northcote and Y.T.J. Kwong, and on contributions by W.R. Bergery of Teck Corporation, L. Bond of Afton Operating Corporation, M. Osatenko of Cominco Ltd. and N. Vollo, Consulting Geologists.

...V.A. Preto

IRON MASK Cu-Au DEPOSITS

Name	Reserve (Production) M short tons	Cu %	Au opt
Afton	Pit 24.4 (1977-87) U.G. 10.5 <i>34.9</i>	0.91 <i>1.50</i> <i>1.08</i>	0.020 <i>0.030</i> <i>.86g</i>
Pothook	2.60 (1987-88)	0.35	0.021
Crescent	1.36	0.46	0.006
Big Onion	Vollo - 2.64 Afton - 3.6	0.84 0.71	0.012 0.012
D.M.	2.96	0.38	0.007
Ajax	27	0.46	0.01
Iron Mask & Erin	0.182 (1904-28)	1.4	?

**Porphyry Cu-Mo-Au
Island Copper Deposit
Vancouver Island, B.C.**

The Island Copper deposit, operated by BHP-Utah Mines Ltd. on northern Vancouver Island, is an island arc-type porphyry Cu-Mo-Au deposit that resulted from the intrusion of a 180Ma rhyodacite dike-like body into comagmatic basalts, andesites and pyroclastics of the mid-Jurassic Bonanza volcanics. Initial estimated ore reserves were 283 million tons of ore at 0.52% Cu and 0.017% Mo. The geology of the deposit has been substantially clarified in recent years. Research studies (O. Arcancibia, Ph.D. thesis in prep.), as well as pit and surface mapping and core logging by company geologists, form the basis for the advances presented herein. Current geological knowledge suggests that the porphyry

Ajax W&E = 24.5 Mt 0.46% Cu, .008 .34g/t Au
Afton pit P&R = 34.9 M tons
31.6 M tonnes @ 1.08% Cu 0.86g/t Au

system evolved from an early, probably juvenile fluid-dominated stage to one strongly influenced by meteoric waters, as the main heat source cooled and further intrusion and brecciation took place. At least three main stages of alteration-mineralization have been differentiated.

1) AN EARLY STAGE, related to the main intrusion of rhyodacite porphyry, involved the development of outwardly progressing zones:

- o a) a stockwork core of quartz-magnetite-amphibole-Na plagioclase;
- o b) a biotite-magnetite zone with chalcopyrite, pyrite and molybdenite;
- o c) a chlorite zone with pyrite and minor chalcopyrite;
- o d) an outermost epidote zone. All are developed in Bonanza Volcanic rocks except the quartz-amphibole-magnetite core, which in addition formed along the margins of the rhyodacite porphyry. The biotite alteration, along with the main copper mineralization, partly overprinted the quartz-magnetite-amphibole core.

2) A STRUCTURALLY CONTROLLED INTERMEDIATE STAGE, superimposed upon the earlier assemblages, was related to the emplacement of a quartz stockwork and is characterized by sericite, chlorite and kaolinite assemblages, with local concentrations of pyrite, molybdenite and chalcopyrite. It is mainly developed in the rhyodacite porphyry and immediate wallrocks. Much of the chlorite overprint currently observed in the pit could have been associated with this stage.

3) A LATE STAGE, related to the emplacement of breccias under extreme base-leaching conditions, is characterized by pyrophyllite, kaolinite, sericite and dumortierite. Further low temperature alteration episodes included "yellow dog" ankerite-calcite veining, widespread zeolite development, and the precipitation of remobilized carbon-bearing organic compounds.

The bulk of copper mineralization was introduced during the early stage, to be followed by the main episode of molybdenum. Almost all of the copper occurs in the form of chalcopyrite, predominantly hosted by biotitized Bonanza Volcanics.

Gold output since production started in 1971 is about 880 000 ounces with an annual rate of slightly higher than 50 000 ounces. This renders Island Copper one of the largest gold producers in British Columbia. Average head-grade in the deposit is about 0.22 ppm Au, with large volumes assaying in excess of 0.4 ppm Au. Only 50% of the gold is recovered in the copper concentrate, which averages 25% Cu and 8 ppm Au. Studies on gold occurrence, gold-contoured values from blast-hole samples, and bulk sampling tests indicate that gold was essentially associated with the early and intermediate stages of alteration-mineralization, although it is not clear whether gold of the latter stage was introduced into the system at that time or remobilized from earlier mineralization. Gold has been observed in the native form, as micron-sized inclusions in chalcopyrite, pyrite, molybdenite and silicates.

Certain features, such as the positive correlation between gold and copper, the overall association of gold with the potassic (biotite-rich) alteration zone, and the high content of magnetite in the system (+8% vol.), are characteristic of

ORE GENESIS RESEARCH

Chemical controls on metal ore grades - an essential part of exploration models for economic ore deposits - are being investigated by the Australian Bureau of Mineral Resources using a new approach adapted from chemical engineering. The first application is on hydrothermal tin-tungsten deposits, where ore fluid compositions are being analysed semi-quantitatively by a new heavy-ion analytical facility (HIAP) proton-microscope technique. The work is being carried out in collaboration with CSIRO's Division of Exploration Geoscience.

most known gold-rich porphyry copper deposits.

Authors: J. Perello, J. Himes - BHP-Utah International Inc; O. Arancibia, P. Clark - Queen's Univ; P. Burt, C. Clarke, C. Fleming & A. Reeves - BHP-Utah Mines Ltd.; and, C. Leitch - U.B.C.

MT. MILLIGAN Alkaline Intrusive Au-Cu Deposit

The Mt. Milligan MBX porphyry gold-copper deposit, 90 miles northwest of Prince George, B.C., has a drill inferred inventory of 100 million tons. Grade ranges from 0.01 to 0.15 oz/t gold and 0.2% to 1.0% copper.

The project is a joint venture between Continental Gold Corp. (70%) and BP Resources Canada Limited (30%). The joint venture holds a large contiguous block of claims covering 42 square miles of the mineral belt, accessible by an all-weather, heavy-duty gravel logging road from MacKenzie, 40 miles to the east.

In 1983 and 1984 BP acquired the property by staking and acquisition and identified a gold-copper soil geochemical anomaly approximately three square miles in extent. In 1985 after IP and magnetic surveys over the geochemical anomaly, BP exposed gold mineralization in trenches on the Creek and Esker zones.

United Lincoln Resources Inc., (a Continental Gold Corp. subsidiary) optioned the property from BP in 1986. Lincoln, having undertaken extensive trenching within the soil geochemical anomaly and additional magnetic surveys, commenced diamond drilling on the Creek and Esker zones and discovered the MBX deposit in 1987.

Regionally the property lies within the central volcanic core of the Upper Triassic-Lower Jurassic Takla Group. Takla volcanics are dominated by subaqueous alkalic pyroxene porphyritic andesite and basalt flows and pyroclastics with subordinate intercalated tuffs and argillites. Intruding the volcanic stratigraphy are comagmatic alkaline syenite, monzonite-diorite stocks.

The MBX porphyry gold deposit is situated on the eastern side of a 3,000 foot diameter porphyritic monzonite stock. Mineralization is hosted by easterly dipping pyroxene porphyritic andesite to latite flows, pyroclastics, interbedded

MINERAL DEPOSITS DIVISION

(MDD) - G.A.C.

Executive & Directors 1989-1990

Chairman: Paul Wojdak

Westmin Resources Ltd.
904-1055 Dunsmuir Street
Vancouver, B.C., V7X 1C4
Tel: (604) 681-2253 FAX: (604) 681-0357

Past-Chairman: Tom Schroeter

B.C. Ministry of Energy, Mines &
Petroleum Resources
159-800 Hornby Street
Vancouver, B.C. V6Z 2C5
Tel: (604) 660-2812 FAX: (604) 660-2653

Vice-Chairman: Ed Debicki

Inco Exploration and Technical Services
Copper Cliff, Ontario, P0M 1N0
Tel: (705) 682-8452 FAX: (705) 682-8243

Program Chairman: Alvin Jackson

Cyprus Metals Canada
1810-1055 West Hastings Street
Vancouver, V6E 2E9
Tel: (604) 685-6867

Secretary: Linda Thorstad

Interaction Resources Ltd.
720 - 800 West Pender Street
Vancouver, B.C., V6C 2V6
Tel: (604) 684-2285 FAX: (604) 684-8887

Treasurer: Robert Cathro

Cathro Exploration Corp.
2320 Bellevue Ave.
West Vancouver, B.C., V7V 1C8
Tel: (604) 926-6003

Publications: Pat Sheahan

Konsult Int. Inc.
44 Gemini Road
Willowdale, Ont., M2K 2G6
Tel: (416) 223-7750 FAX: (416) 361-0369

Newsletter - The Gangue: Brian Grant

B.C. Ministry of Energy, Mines &
Petroleum Resources
525 Superior Street, Rm 122-10
Victoria, B.C., V8V 1X4
Tel: (604) 356-1693 FAX: (604) 356-8153

MDD Directors:

Robert Hewton Vancouver	(604) 687-2557
Alex Davidson Vancouver	(604) 681-3771
Lee Pigage Whitehorse	(403) 668-3578
Randy Maass Timmins	(705) 268-6644
William Pearson Toronto	(416) 368-4636
Ken Shannon Toronto	(416) 947-9166
David Moore Toronto	(416) 869-1850
Jacques Trottier Montreal	(514) 393-1505
Geoff Thurlow Corner Brook	(709) 639-7931

trachytic tuffs and a 30 to 150 foot thick porphyritic monzonite dike. Mineralization persists into the eastern margin of the monzonite stock but with decreasing grades.

The multi-phase monzonite stock is fine to medium grained with 20% plagioclase laths in an aphanitic potassium feldspar matrix. This is one of three alkaline plutons on the property that form a north-northeast alignment, suggesting a structurally controlled emplacement. The porphyritic character of the intrusive units and the number of dikes is indicative of a hypabyssal subvolcanic environment.

Potassium silicate and propylitic alteration assemblages have formed outwardly from the stock for 1000 and 8000 feet respectively. An early, fine grained felted hydrothermal biotite superimposed on the volcanics is over printed by pervasive grey potassium feldspar. Biotite comprises 10 to 35 per cent of the volcanic units and potassium feldspar up to 50 per cent, while pyroxene is typically replaced by actinolite within the potassic zone. In fine grained laminated tuffs massive potassium feldspar replacement is common. The potassium silicates overprint the enclosing propylitic assemblage which is principally comprised of epidote, carbonate and pyrite. Veinlets of magnetite with minor chalcopyrite post date the main period of sulphide precipitation and are restricted to the potassic zone. Weak but pervasive sericite partially replaces plagioclase phenocrysts in the monzonite stock and in the porphyritic monzonite dike within the zone of potassic alteration. Throughout the MBX deposit, alteration is typically pervasive and veining is infrequent.

Disseminated grains and coalescing grain aggregates of chalcopyrite and pyrite comprise approximately 60% of the total sulphide content. Fracture controlled pyrite-chalcopyrite mixed veinlets are less abundant. Chalcopyrite and pyrite-bearing K-feldspar-carbonate veinlets are relatively rare. Quartz veining is noticeably absent. The distribution of sulphides is zoned but not uniformly. At the north end of the MBX deposit pyrite and chalcopyrite occur in equal concentrations. Bornite, though present, is a minor constituent. Within the west central portion of the deposit the pyrite:chalcopyrite ratio is approximately 3:1. Along the east and southeastern margin of the deposit the pyrite content increases to 5 to 10%, and the pyrite:chalcopyrite ratio is approximately 20:1.

Metal zoning closely parallels that of the sulphides with gold concentrations increasing with the pyrite:chalcopyrite ratio. This relative gold enrichment, coincides with the transition from potassium silicate to the propylitic alteration assemblage.

Both chalcopyrite and pyrite are auriferous and occur as separate grains. Intergrown sulphides are rare, an important positive metallurgical feature. Gold associated with pyrite occurs as small particles on grain margins.

At the West Breccia zone, situated on the western flank of the monzonite stock, an intrusive breccia, comprised mostly of monzonite and a few scattered volcanic clastics, is enclosed by a broad zone of brecciated and fractured monzonite. Within the intrusive breccia pink secondary potassium feldspar is pervasive, however, moving outward from the breccia, the potassium feldspar is fracture controlled. Both chalcopyrite and pyrite occur as fracture fillings and as discrete fine grains partially replacing altered mafic minerals. Chalcopyrite predominates over pyrite and gold concentrations are low.

The Creek, Esker and 79 zones are located within propylitically hornblende-pyroxene porphyritic andesites approximately 1000 feet southwest and 1500 feet west, respectively of the MBX porphyry deposit. These zones are northeast striking, steeply northwest-dipping, tabular bodies of auriferous semimassive to massive sulphides occupying fracture systems radial to the monzonite stock. Each of the deposits comprises three to five subparallel, sulphide-rich bodies, spaced across 200 to 300 feet. Individual structures range from 1 to 10 feet in thickness and grade from 0.10 to 2.89 oz/ton gold and 0.2 to 10% copper. Each zone is open along strike and down dip. The propylitically altered andesites between the individual sulphide bodies carry 30 to 350 ppb gold and 200 to 2500 ppm copper. Silver, arsenic and antimony are present at background concentrations.

Pronounced metal zonation is related to the proximity to the monzonite stock. The MBX porphyry deposit adjacent to the stock, contains less than 10 ppm combined arsenic and antimony. At the Creek, Esker and 79 zones, combined concentrations increase to 300 ppm and to 2000 ppm respectively. Silver shows a similar relationship with concentrations ranging from 1.5 ppm in the MBX zone, to 10 to 40 ppm in the Creek zone, and 60 to 200 ppm in the Esker zone.

Mark Rebaghatti, P.Eng., who conceived the original exploration program and Dave Copeland, P.Eng., are Vancouver-based consulting geological engineers who have directed and managed this project to its current stage.

[Ed. Note: Recent news releases from Continental Gold Corp. indicate the Mt. Milligan Cu-Au deposit is a massive, blanket-shaped deposit 4,300 feet long, up to 3,100 feet wide and up to 800 feet thick. Over 150,000 feet of drilling has been completed in 200 holes. Reserves are reported to be in the 150 to 200 million ton range, grading 0.3% copper and 0.02 oz/ton gold].

STEWART - ISKUK BELT Northwestern B.C.

K-Feldspar Connection: Relationship of K-Feldspar Intrusions to Cu Porphyries and Au Veins, Stewart-Iskut Belt

Hypabyssal copper-gold porphyries and mesothermal to epithermal gold-silver veins in the Stewart-Iskut gold belt are related to Early Jurassic intrusive rocks which are either alkaline or calc-alkaline in composition. In the Premier and Sulphurets areas the intrusive rocks feed and cut a Lower Jurassic volcano-sedimentary sequence of the Hazelton Group. At the Snip, Skyline and Galore Creek deposits the intrusive rocks cut Upper Triassic, Stuhini Group volcanic and sedimentary rocks. Porphyry and vein-type ore deposits are linked by:

- 1) their spatial and inferred temporal association with intrusive rocks, in particular distinctive two-feldspar porphyry with K-feldspar megacrysts (Premier porphyry is a well documented example)
- 2) potassic alteration - pervasive sericite, K-feldspar (adularia), and less commonly biotite
- 3) a structural control, interpreted to be syn-volcanic (ex.

Premier and Galore Creek)

At Premier, hornblende-plagioclase-K-feldspar-quartz porphyritic dacite dikes (Premier porphyry), derived from the Texas Creek granodiorite batholith, are emplaced along intersecting northeast and northwest structures. These appear to control volcanic stratigraphy and are interpreted to be synvolcanic faults. The ore consists of quartz-K-feldspar (adularia) veins, stockwork and breccia that follow the same structures as Premier porphyry. Past production at premier is 4.7 Mt at 2.2 g/t Au and 80.3 g/t Ag. Mineralization dies out abruptly near overlying maroon (oxidized) strata, which suggests that ore deposition resulted from mixing of hydrothermal fluid with meteoric water. Silver and gold are most abundant in the upper part of the deposit, abundance of base metals and mineral grain size increase at depth. The sericite alteration zone flares outward toward the top of the deposit. These are classic features of an adularia-sericite epithermal deposit. Gold occurs primarily as electrum that has a close spatial association with tetrahedrite.

Mineralization in the Sulphurets area is controlled by two north-trending structures; the steep Brucejack fault and the Sulphurets fault, whose shallow west dip results in a sinuous map pattern. These, and secondary structures, control Early Jurassic diorite, syenite and granitic intrusions and localize sericite-K-feldspar alteration. An alteration zone of this type hosts the West Zone deposit (0.775 Mt at 12.1 g/t Au and 786 g/t Ag) adjacent to the Brucejack fault. Mineralization is an anastomosing network of vein stockworks and breccia, interpreted to be a silicified shear zone, bounded by intrusive K-feldspar porphyry.

The Kerr deposit (60 Mt at 0.84% Cu, 0.34 g/t Au and 2.05 g/t Ag) occurs within sheared and sericite-altered volcanic rocks that are bounded by splays of the Sulphurets fault. A diorite body, K-feldspar porphyry dikes and high grade Au-Ag-Cu veins are all contained within the north-trending structural zone. The Kerr porphyry deposit contains pyrite, chalcopyrite, tennantite, bornite and chalcocite occurring as disseminations and disrupted quartz veins.

The Snip deposit (Twin zone 1.43 Mt at 21.9 g/t Au) is a mineralized shear zone that trends 120 and dips moderately to the southwest (about 55). Discordant mineralization, that cuts massive feldspathic wackes, comprises pyrite, pyrrhotite, chalcopyrite, sphalerite, galena and arsenopyrite. The ore is both massive and strongly foliated, quartz is commonly brecciated with a crackle texture. Mineralization is restricted to the shear zone and contacts with wallrock are sharp. Potassic alteration is represented by pervasive biotite-flooding and irregular K-feldspar (adularia) replacement of the wallrock.

The Johnny Mountain gold mine, 1 km southeast of Snip, is a structurally disrupted mesothermal gold-bearing quartz vein deposit. Current reserves are 0.622 Mt at 19.5 g/t Au and 0.75% Cu. Silicification and potassic alteration occur along a series of northeast-trending structures in close proximity to an orthoclase porphyry.

The Galore Creek alkalic porphyry Cu-Au deposit (125 Mt at 1.06% Cu, 0.40 g/t Au and 7.7 g/t Ag) comprises 10 tabular to manto-shaped deposits. Mineralization is hosted in volcanics, breccia pipes and K-feldspar megacrystic syenite dikes and plugs. The mineralization and several phases of syenite porphyries are in part controlled by north and northeast

structures, interpreted to be synvolcanic. The deposits are characterized by skarn-type, metasomatic or late magmatic alteration products (epidote, garnet, diopside and magnetite) along with pervasive K-feldspar and biotite alteration, anhydrite and gypsum. Ore minerals consist of pyrite, chalcocopyrite, magnetite, bornite and minor sphalerite and galena.

Porphyry and mesothermal mineralization could be incorporated into the Premier ore depositional model. This would suggest synchronous mineralization and depth/temperature dependency in a single stage process. This is probably too simplistic. Intrusive, structural and mineralization histories are known to be multi-stage and the ore fluids probably range from magmatic to mixed juvenile/meteoric. More work needs to be done to answer questions concerning:

- 1) age, displacement, history of ore controlling fluids
- 2) relative and absolute ages of mineralization types
- 3) composition of fluids

The authors acknowledge informative discussions with: R. Britten (Esso Minerals); F. Hewitt (Northair Group); B. Hewton and B. Butterworth (Sulphurets Gold Corporation); R. Nichols, I. Paterson and B. Wolfe (Cominco Ltd.)

...Derek A. Brown
Paul Wojdak

FUTURE MEETINGS, WORKSHOPS & FIELDTRIPS

OCTOBER 1989

14 - 19th **STRATIGRAPHY AND METALLOGENY OF THE SICKER GROUP, VANCOUVER ISLAND.** Workshop and fieldtrips, Nanaimo, B.C. Details: H. Paul Wilton or Nick Massey, c/o B.C. Geological Survey Branch, 220-756 Fort Street, Victoria, B.C., Canada, V8W 3A3. Telephone (604) 356-2818.

NOVEMBER 1989

5 - 8th **WORLD GOLD '89**, Reno, Nevada. Details: Meetings Dept. SME, Box 625002, Littleton, Colorado, USA 80162, Tel. (303) 973-9550.

20 - 21th **MODERN EXPLORATION TECHNIQUES.** Conference sponsored by the Saskatchewan Geological Society. c/o Robert Troyer, SGS Symposium '89, P.O. Box 234, Regina, Saskatchewan, S4P 2Z6. Tel: (306) 787-2562.

DECEMBER 1989

3-5th **ORE MICROSCOPY AND FLUID INCLUSION ANALYSIS SHORT COURSE**, Pullman Washington. Registration Fee \$375 U.S. Contact: Dr. Lawrence D. Meinert, Ore Microscopy Short Course, Washington State University, Pullman, WA 99164-2812.

6 - 8th **NORTHWEST MINING ASSOCIATION 1989 ANNUAL MEETING**, Spokane Washington. One of the

highpoints of the mining year! Details: NWMA, 414 Peyton Bldg., Spokane WA 99201, Tel. (509) 624-1158.

FEBRUARY 1990

6 - 9th **CORDILLERAN ROUNDUP.** This is the premier exploration and mining show in western Canada. Vancouver Hotel, Vancouver, B.C. Details: Jack Patterson, Manager, B.C. & Yukon Chamber of Mines, 840 W. Hastings St., Vancouver. Tel: (604) 681-5328.

MARCH 1990

1 to 11th. **MEXICAN SILVER DEPOSITS EXCURSION, 1990.** Sponsored by Society of Economic Geologists (SEG) in conjunction with SME/AIME Annual Meeting in Salt Lake City, Utah, Feb. 26 - Mar. 1, 1990. Est. cost US\$850. Limit 35 participants. Reservations & Information contact: Kenneth F. Clark, Economic Geology Publishing Co., P.O. Box 637, University of Texas, El Paso, El Paso, Texas, 79968-0637, USA. Tel: (915) 533-1966, FAX: (915) 544-7416.

9 - 10th **CAME SYMPOSIUM - Computer treatment of Exploration and Mining Data - Do's and Don'ts.** Fee Cdn\$125. Details: P&D Assoc. of Canada, Suite 1002, 74 Victoria Street, Toronto, Ont., M5C 2A5. Tel: (416) 362-1969 FAX (416) 362-0101.

9 - 11th **EXPLORATION GEOCHEMISTRY: Design and Interpretation of Soil Surveys.** Fee Cdn\$400. Metro Toronto Convention Centre. Sponsored by the Association of Exploration Geochemists and the Prospectors and Developers Association of Canada. Contact PDAC: Tel:(416) 362-1969 or FAX: (416) 362-0101

11 - 14th **PROSPECTORS & DEVELOPERS ASSOCIATION Annual Convention.** Theme: Exploration & Development - New Realities. Royal York Hotel, Toronto, Ont., Canada. Contact PDAC: Tel:(416) 362-1969 or FAX: (416) 362-0101.

APRIL 1990

1 - 5th **GEOLOGY AND ORE DEPOSITS OF THE GREAT BASIN**, symposium and field trips, Sparks, Nevada. Details: Geol. Soc. Nevada, Box 12021, Reno, NV., USA 89510, Tel. (702) 786-0870.

MAY 1990

May 14 -15 **FLUIDS IN TECTONICALLY ACTIVE REGIMES OF THE CONTINENTAL CRUST.** Hotel Vancouver, Vancouver, B.C., Preceding 1990 Annual GAC/MAC Convention. Tentative fees \$250 (\$150 for students). Contact: Dr. B.E. Nesbitt, Dept. of Geology, Univ. of Alberta, Edmonton, T6G 2E3. Tel: (403) 492-5071, FAX (403) 492-2030.ts

May 28 - June 1 **CIMM Symposium on the POLYMETALLIC BELT OF NORTHWESTERN QUEBEC**, Rouyn-Noranda. Details: Maurice Rive, Ministre de l'Energie et des Ressources, 19 rue Perreault Ouest,