

DETAILED BEDROCK  
GEOLOGY OF THE BRENDA  
COPPER-MOLYBDENUM MINE,  
PEACHLAND, BRITISH COLUMBIA

802410

by

WILLIAM MICHAEL ORJEL

M. Sc. THESIS, U. B. C

924/16

MAY, 1972.

**DETAILED BEDROCK GEOLOGY  
OF THE BRENDA COPPER-MOLYBDENUM MINE  
Peachland, British Columbia**

by

**WILLIAM MICHAEL ORIEL  
B.Sc., 1968  
Colorado State University**

**A THESIS SUBMITTED IN PARTIAL FULFILMENT OF  
THE REQUIREMENTS FOR THE DEGREE OF  
MASTER OF SCIENCE**

**in the Department  
of  
Geology**

**We accept this thesis as conforming to the  
required standard**

*G. J. Sinclair*.....  
*Delavault*.....  
*[Signature]*.....

**THE UNIVERSITY OF BRITISH COLUMBIA**

**May, 1972**

## ABSTRACT

Detailed mapping of the bedrock geology of the Brenda copper-molybdenum deposit has revealed a consistent sequence of geological events. Consolidation of the quartz diorite host rock that forms Brenda stock was followed by intrusion of small pegmatite and aplite dikes that probably relate to a residual fraction of the parent magma. Age dating of various quartz diorites (White and Harakal, unpublished) indicates that magma solidification occurred about 176 m.y. A concentration of biotite model ages at about 146 m.y. seems best interpreted as the time of ore mineralization.

Mineralization occurred at the beginning of a continuous sequence of related structural events. All mineralized ore fractures and younger shears, faults and intramineral dikes exhibit similar, strongly preferred orientations that are unlike orientations of felsic dikes or primary foliations in the quartz diorite host rock.

The ore occurs entirely in fractures as products of three sequential overlapping phases of mineralization and consists almost entirely of the ore minerals chalcopyrite and molybdenite with quartz gangue predominanting.

Hydrothermal alteration products exist mostly as thin envelopes of potassium feldspar and/or biotite and propylitic alteration. Large argillic alteration zones also exist, however, and are associated commonly with extensively sheared zones. An effort was made at finding a zoning pattern of ore minerals, minor elements and alteration minerals. Ore mineral zoning in the classical sense appears to be absent at Brenda. Metal grades show a crude zonal distribution

that correlates directly with density of ore fractures. hydrothermal alteration minerals and minor elements in chalcopyrite also lack any obvious systematic zonation pattern on the scale of the mine pit.


Intramineral dikes of various textures and compositions have been dated at about 130 m.y. They formed after ore mineralization yet before late minor veining not associated with ore-producing veins.

A. J. S.

## TABLE OF CONTENTS

CHAPTER		PAGE
I	INTRODUCTION	1
	PURPOSE	1
	PHYSIOGRAPHY	1
	HISTORY	3
	METHODS OF STUDY	5
II	GENERAL GEOLOGY	8
	INTRODUCTION	8
	NICOLA GROUP	8
	TERTIARY (?) CONGLOMERATE	10
	INTRUSIVE ROCKS	10
III	GEOLOGY OF BRENDA MINE - STRUCTURE	15
	FELSIC DIKES	15
	INTRAMINERAL DIKES	22
	ZONES OF SHEARING	23
	MOLY-SLIPS	25
	BLACK CLAYS	25
	FAULTS	27
	JOINTS	29
	AGE DATING	29
	SUMMARY	34
IV	GEOLOGY OF BRENDA MINE - VEIN DESCRIPTIONS	38
	INTRODUCTION	38
	STAGE I VEINS	40
	MINERALOGY	40
	STRUCTURE	42
	STAGE II VEINS	51
	MINERALOGY	51
	STRUCTURE	54
	STAGE III VEINS	55
	MINERALOGY	55
	STRUCTURE	57
	STAGE IV VEINS	58
	STRUCTURE	61
	SECONDARY MINERALS	61
	HYDROTHERMAL ALTERATION	67
	SUMMARY	71

CHAPTER		PAGE
V	ZONING	74
	MINERAL ZONING	74
	COPPER AND MOLYBDENUM	74
	LEAD	77
	MINOR ELEMENT ZONING	81
VI	SUMMARY AND CONCLUSIONS	83
	GENERAL GEOLOGICAL EVOLUTION OF THE BRENDA DEPOSIT	83
	UNUSUAL ASPECTS OF THE BRENDA DEPOSIT PROBLEMS FOR FUTURE INVESTIGATION	85 87
	BIBLIOGRAPHY	89
	APPENDIXES	91
	INDUCED POLARIZATION	91
	SOIL GEOCHEMISTRY	91



## CHAPTER II

## GENERAL GEOLOGY

INTRODUCTION

Small scale geologic mapping surrounding the Brenda Mine was done by J. M. Carr in 1967 for the B. C. Department of Mines (Figure 2). It is not the purpose of this thesis to expound further on the regional geology, hence Carr's study serves as the main source of regional geologic information. There are two main rock types: the Nicola Group tuffs and argillites and the intrusive quartz diorites that contain the Brenda ore body.

NICOLA GROUP

Carr (1967) defines four within the Nicola Group units that crop out west of Brenda stock. These are porphyritic tuff, schistose hornfels, greywacke, and argillite-limestone sequence. The schistose hornfels lies in contact with intrusive rocks for one mile south and north of the orebody and has an outcrop zone approximately 1/4 of a mile wide. Megascopically, it is a fine-grained, silica-rich rock possessing foliation cleavage and lineations as a result of a concentrated development of biotite plates up to four millimeters wide. Petrographically, it is a "fine-grained, quartz-feldspathic mosaic containing disseminated biotite and chlorite, lenticular knots of biotite, single unzoned plagioclase crystals and locally, grains of quartz" (Carr, 1967, p. 186). Schistosity strikes consistently N12°W and dips 40°SW. Probable original



bedding in the tuffs around Brenda Lake strike N80°W and dip 65°S. The hornfels probably is a metamorphic equivalent of the porphyritic tuff and tuff breccia units recognized by Carr.

Porphyritic tuffs crop out in a wide northerly pinching zone closely paralleling the schistose hornfels. Exposed width of the outcrop zone at the southern end of the unit exceeds one mile. Contact relations with argillites and graywackes are mostly hidden by deep glacial drift. They contain abundant fragments of volcanic material such as porphyritic andesites and dacites. Volcanic flows were not mapped in the region, however. Carr describes tuffs in the eastern part of the unit as a tuff breccia containing rounded to subangular volcanic rock fragments up to two inches long. In the western half the tuffs are characterized by massive, fine-grained or glassy rock interlayered or intermixed with clastic sedimentary material (Carr, 1967).

Graphitic argillite and limestone units (numbers 5 and 4 respectively on Carr's map) crop out about two miles west of the mine over an area extending for two miles north and a mile south of the pit. They consist of two black limestone beds 200-300 feet thick surrounded and separated by tuffaceous greywackes, siltstones and graphitic argillites. "The limestones are graphitic, recrystallized, schistose, fine-grained rocks containing rare quartz grains and particles of chert. Their schistosity shows minor drag-folds plunging southward at 40 degrees." The black argillites and greywackes "are thin-bedded, incompetent rocks, and the soft graphitic argillite has in places squeezed discordantly into faults." (Carr, p.185).

Another type of Nicola rock in the area is the greywacke unit. These are bedded volcanic greywackes and siltstones that apparently grade into increasingly metamorphosed hornfels close to the intrusive contact. Scour features in the siltstones suggest to Carr that the beds are upright. The hornfels are brown, quartzitic rocks with large amounts of finely dispersed biotite. Sericite was noted on cleavage surfaces. Fine-grained quartz, plagioclase, biotite and hornblende are the dominant constituents. Irregular patches of pale and/or dark minerals are common giving the rock a somewhat mottled appearance.

#### TERTIARY (?) CONGLOMERATE

A conglomerate of possible Tertiary age (Carr, 1967) is located one mile north of Brenda Lake on the old Noranda road. It consists of rounded pebbles and cobbles in a fine-grained matrix; all possessing an irregular, black manganiferous coating. The rock is poorly sorted with fragments of tuff, chert, quartz diorite and porphyry in an argillaceous matrix. It was also reported to have been glacially striated; however, when the writer examined the exposures, all traces of glacial polishing had been removed through bulldozer road clearing.

#### INTRUSIVE ROCKS

The Brenda ore body is within a body of quartz diorites termed by Carr the Brenda Stock. It appears more linear than equant, extending in a northerly direction. To the northwest, north and east, however, its limits have not been defined. The stock has a

minimum width of 4 miles east to west and a length greater than 7 miles. To the immediate south is the Similkameen batholith and to the northwest is the Pennask batholith. The Brenda vicinity has been included as part of both the Similkameen and the Pennask batholiths by Peto (1970) and Schau (1968) respectively. Further work is clearly needed. The western contact of the intrusive is with Nicola tuffs and argillites as described previously. Nearly 3 miles north of the mine the contact diverts from a northerly to northwesterly trend.

Four successive subparallel quartz diorite units and a later intrusive phase were defined by Carr as comprising the Brenda Stock. These are the Medium, Speckled, Uniform, Porphyritic, and Fine quartz diorites. Separation of units was based on macroscopic textural differences arising primarily from small changes in the distribution, size, and ratios of biotite and hornblende. Basically, all units are similar; they are all quartz diorites possessing slightly varying amounts of quartz, plagioclase, potash, feldspar, hornblende, biotite, and magnetite, sphene and apatite; the latter three occur as accessory minerals. Quartz forms about 25 percent, plagioclase 50 percent, orthoclase and microcline up to 20 percent; and hornblende and biotite account for 10 to 30 percent of the rock volume.

Many outcrops possess a faint foliation due to the alignment of mafic minerals, especially biotite. Foliation attitudes are generally parallel to the contact of the adjacent unit of the Nicola contact. Most contacts dip from vertical to steeply westward.

"Medium Quartz Diorite(7)....A rough-weathering rock, the Medium quartz diorite is relatively dark and inhomogeneous, although less so with distance from the outer contact, and it has a medium-grained appearance due to a high content of crystals larger than 2 millimetres. Foliation is mostly well developed, and the rock is commonly streaky due to local variation of the dark mineral content. Hornblende mostly exceeds biotite in amount, and its crystals may reach a length of 2 centimetres. Quartz is partly wedge-shaped, and the larger grains have a blue colour in some outcrops. Potash feldspar is less abundant in this rock than others. The estimated range of modal compositions of the Medium quartz diorite is as follows: Quartz, 10 to 30 percent; potash feldspar, 0 to 5 percent; plagioclase, 45 to 60 percent; hornblende and biotite, 10 to 30 percent; magnetite, etc., 1 percent."

"Speckled Quartz Diorite(8)....Characteristic features aiding recognition of the rock include: A speckled appearance due to the interspersions of small biotite and hornblende crystals with larger ones; the presence of biotite in amounts equalling or exceeding those of hornblende, and its occurrence chiefly in sieve-like shapeless plates of size to 4 millimetres; a prismatic or needle-like shape of most hornblende crystals; and a finely granular appearance of most quartz. Quartz tends to be blue-coloured, as in certain other units. The estimated range of modal compositions of the Speckled quartz diorite is as follows: Quartz, 20 to 25 percent; potash feldspar, 10 to 15 percent; plagioclase, 45 to 55 percent; hornblende and biotite, 15 percent; magnetite, etc., 1 to 2 percent."

"Uniform Quartz Diorite(9)....Like the Speckled rock, the Uniform quartz diorite is predominantly medium grey in colour. It is distinguished chiefly by the even distribution of its dark minerals, of which hornblende is the best shaped and most abundant. Biotite occurs mostly in small plates, which are partly euhedral, and rarely in books as large as 3 millimetres. Quartz is also mostly fine grained but locally makes aggregates as large as one-half centimetre and containing small feldspars. The estimated range of modal composition of the rock is as follows: Quartz, 20 to 30 percent; plagioclase, 40 to 55 percent; potash feldspar, 5 to 15 percent; hornblende and biotite, 12 to 15 percent; rest, 1 percent."

"Porphyritic Quartz Diorite(10)....The rock is lighter coloured than others, and it generally has few inclusions. Dark minerals are relatively sparse and scattered and, together with quartz which forms 1/2-centimetre aggregates, they give the rock a porphyritic appearance. Biotite is typically in well-shaped books of size to one-half centimetre and it exceeds hornblende in amount. Hornblende makes relatively few crystals as large as 2 millimetres; quartz fails to occur in wedge shapes. The estimated range in mode of the rock is as follows: Quartz, 20 to 30 percent; plagioclase, 45 to 60 percent; potash feldspar, 8 to 15 percent; biotite and hornblende, 10 to 15 percent; rest, 1 to 2 percent."

"Fine Quartz Diorite(11)....Although appearing rather like the Speckled quartz diorite, the Fine quartz diorite is somewhat finer grained and lighter coloured; it possesses quartz partly as small wedges, hornblende partly in excess of biotite, and biotite in plates and thick books mostly smaller than those in the Speckled quartz diorite. Additional useful features for identification are blue quartz grains, a well-developed foliation, unusually abundant sieve-crystals of potash feldspar, and well-shaped hornblendes and biotites distributed patchily in the rock." (Carr, 1967, pp. 190-191)

It should be noted here that the writer collected more than 70 samples of the above mentioned quartz diorites from random outcrops outside the pit area for a radius of two miles. Without predetermination, the writer grouped these samples into macroscopic textural divisions which conformed to Carr's as closely as possible. It was found that there is little correlation between a mapped unit and the textural descriptions of the rocks that the writer collected within its boundaries. Each unit as shown on Carr's map contains more rock types than the name applied to the unit. In several locations, especially in the porphyritic unit, there were as many as three different rock types from the same outcrop. The writer noted that if foliation is well-developed the texture of any particular sample can vary considerably depending on whether it is broken parallel or perpendicular to the foliation. Variable character of weathered surfaces can produce superficial differences in rock type that can lead to ambiguity in applying Carr's classification. Although Carr's regional mapping has provided much needed information and insight to the problems, the writer feels that further detailed work is required.

Other separate and possibly distinct quartz diorite bodies crop out in the area. A "Southern Quartz Diorite" occurs in the southwest corner of Carr's map about 2 miles below the pit. It is

similar in appearance to the aforementioned units, but Carr suggests that the hornblende-biotite ratio is slightly different and that hornblende does not have the needle-like appearance commonly found in the units of the Brenda Stock. Ground cover obscures contact relations.

The Satellite quartz diorite is a fine-grained distinctively grey, small plug isolated in Nicola greywackes 2 miles north of the mine. Scattered small (1/2 cm) phenocrysts of plagioclase, hornblende and biotite appear in this unit. It also has a well-defined foliation.

Quartz monzonites appear in the southeastern part of Carr's map area and are similar in appearance to the quartz diorite units except that they contain more potash feldspar than the quartz diorites. These rock types are described more thoroughly in Carr's report to which the reader is referred.

Dikes in the region vary from aplites to basalts with several intermediate varieties. Most strike northwesterly and dip steeply. Carr dates some as being pre-ore and other as post-ore. He describes several structural belts as being defined by the presence or absence of aplite-pegmatite dikes of pre-ore age. Detailed data collected by the writer in and around the pit do not support these conclusions. A more detailed discussion of the dikes is presented in the following chapter.

A suggested age of the quartz diorite from unpublished potassium-argon age dating data (Harakal and White, personal communication, 1972) is about 176 million years. A detailed presentation and analysis of the data is located in the following chapter.

## CHAPTER III

## GEOLOGY OF BRENDA MINE - STRUCTURE

Primary structure features of the Speckled quartz diorite that contains the Brenda ore deposit have been referred to in an earlier section. Secondary structural features in their approximate order of relative age are from oldest to youngest: ore veins, intramineral dikes, shear zones, moly-slips, and joints. Ore veins are discussed in detail in a subsequent chapter.

FELSIC DIKES

Felsic dikes appear throughout the pit. They are divided for convenience into three textural divisions: aplite, pegmatite and granophyre dikes. As a group, they represent the earliest recognized geological event subsequent to development of the Speckled quartz diorite. Dikes range in width from 1/4 inch to a maximum of 12 inches. Wider dikes are more persistent laterally than are small dikes, but observed strike lengths do not exceed one hundred feet. The dikes commonly occur in small swarms resembling filled, en echelon, tension fractures. Most dikes are a pale to medium pink color; some, however, are grey. Staining revealed that grey dikes contain less potash feldspar than do pink dikes. (Figure 4)

Aplites have a "sugary texture and consist of quartz, microcline, plagioclase, and minor amounts of green biotite with average grain diameters ranging from 0.05 to 0.3 millimeters....".

however, that mineralization was the agent that reset some biotite and precipitated hydrothermal biotite related to mineralization. If this were so, then the maximum age of mineralization is about 146 m.y. and could be as low as 130 m.y. if separate biotites were mixed for analyses.

The writer prefers the second hypothesis because the first hypothesis appears to be ruled out by all the data so far collected. And the third hypothesis requires extreme coincidence that several random samples would all contain the same relative proportions of separate biotites.

#### SUMMARY

The time of emplacement of the Speckled quartz diorite appears to be about 176 million years ago or lower Jurassic time as indicated by potassium-argon ages on hornblende by Dr. W. H. White of the University of British Columbia (Harakal and White, personal communication, 1972). This intrusion was followed by the introduction of aplite, pegmatite and granophyre dikes possibly related to late magmatic fluids resulting from cooling crystallization and settling of the magma. They strike predominantly northwest and dip moderately to the southeast. Their orientation is significantly different than all other secondary structures, which leads the writer to believe that they are not contemporaneous with or even caused by the same conditions that formed later events as suggested by Carr (1967). Following these after some indeterminate time are the ore veins, intramineral dikes, shears and faults forming in overlapping sequence (Figure 11). Age dating indicates that



mineralization probably commenced about 146 m.y.

Intramineral dikes are termed such for their occurrence in time with Stage IV veins and shears. They are overlapped by both but occur well after the formation of the ore veins. Orientations of intramineral dikes are remarkably consistent striking N59°W, dipping 89°SW; however, their compositions range between andesitic and basaltic. A few trachyte dikes are present that have been described previously by Carr (1967). Two dikes of andesitic composition are apparently of premineral origin. Their relation to the other dikes are unknown. The actual time of the sequence of alteration, mineralization and the formation of dikes and shears cannot as yet be determined absolutely. Two age determinations of mafic dikes have been carried out (Harakal and White, personal communication, 1972) and indicate an age of 130 million years. It appears possible that these dikes, though late in the sequence represent the final events with negligible but significant mineralization following emplacement of intramineral dikes.

Large shear zones averaging about 10 feet wide are predominant in the pit. Shears overlap the intrusion of the intramineral dikes and the end of the period of ore mineralization. Stage IV veins and a few Stage III veins in part intersect the shear zones. Argillic alteration has greatly extended the limits of shear zones and effectively weakens the rock for percolation of ground water and further decomposition of the rock.

Many faults are related to shearing whereas others appear unrelated. Movement direction and ages are difficult to determine and little can be said of them. Faults probably occur contemporaneously

## Structural History at Brenda




Figure 11. Sequence of events showing progressive overlap and relative ages. Stipled area indicates time gap.

with and/or slightly later than shear zones. Some faulting might be of Tertiary age. Orientation of both shears and faults are similar, striking about  $N68^{\circ}E$  and dipping  $75^{\circ}$  to  $85^{\circ}$  southeast.

One of the most important conclusions is the apparent significance of the continuity of orientations of structural features of various ages. This suggests that the events were all in response to the same stress environment that existed with intermittent release over a protracted time period.

Joints are unmineralized late features possibly related to uplift and erosion. Their predominant orientation is  $N80^{\circ}W/25^{\circ}N$ . By comparing the orientation of these features (including mineralized veins discussed in the following chapter), a pattern can be observed with two predominant attitudes  $N60-68^{\circ}E/75-85^{\circ}SE$  and  $N60^{\circ}$  to  $70^{\circ}W/80-85^{\circ}SW$ . These orientations appear to be consistent throughout the pit barring gaps and slight differences due to inadequate exposure and few recorded observations.

## CHAPTER IV

## GEOLOGY OF BRENDA MINE - VEIN DESCRIPTIONS

INTRODUCTION

Chalcopyrite and molybdenite are the abundant ore minerals in the Brenda ore deposit. They occur in three paragenetically and mineralogically distinct vein systems. A fourth stage of veins are late and contain only infrequent sulfide occurrences.

	<u>Time</u>	<u>Primary Opaques</u>	<u>Dominant Gangue</u>
Ore veins	Stage I Veins	Chalcopyrite - Py	Biotite
	Stage II Veins	Cpy-Moly-Py	Quartz K-felspar
	Stage III Veins	Moly-Cpy-Py-Mag-Hem	Quartz Calcite
	Stage IV Veins	(Py, Spec, Hem, Cpy)	Various

Previously Soregaroli (1971), mapping within the pit in 1968, divided the veins into four types: 1) quartz-potash, feldspar-sulfide, 2) biotite-chalcopyrite, 3) quartz-molybdenite, and 4) epidote-magnetite-molybdenite veins, occurring in that order formation. The writer departs somewhat from this order. Biotite veins containing chalcopyrite are found to be consistently separated in time from barren biotite veins that crosscut all other veins except late calcite and epidote veins. Specific vein orientations are slightly more emphasized or de-emphasized by the writer than by Soregaroli. These are discussed in their appropriate order. Much of this work serves to confirm the excellent geology and mineralogy by Soregaroli with much fewer exposures than were available to the writer.

Average grade for the entire pit is 0.18 percent copper and 0.049 percent molybdenum, these metals being present almost entirely as chalcopyrite and molybdenite respectively. Computer calculated reserves are 177 million tons. Gold and silver assays in the copper concentrate average 0.056 oz./ton and 3.72 oz./ton respectively (D. F. Whitford, Brenda geologist, personal communication). Rhenium content in molybdenum concentrate ranges from 120 to 180 ppm (A. E. Soregaroli, personal communication). Ore minerals occur almost entirely in fractures, and fracture density is the dominant control of ore grade. Disseminations occur only rarely in areas where alteration is intense.

Other hypogene ore minerals are pyrite, magnetite, hematite, bornite, sphalerite, and galena. Galena has thus far not been observed in place or in polished sections by the writer. Sphalerite and bornite are rarely visible macroscopically but are in places apparent in polished sections examined by reflected light microscopy. Pyrite averages about one percent of vein sulfides. Amounts of the other ore minerals mentioned vary drastically from place to place but normally are found only in trace amounts.

Gangue minerals include quartz, potassium feldspar, calcite and clinozoisite. Barite was found in one location in the pit. Occurrences of the above minerals are discussed in detail in following paragraphs.

## STAGE I VEINS

The earliest sulfide minerals in the intrusion were deposited in biotite-lined fractures, 1/32 to 1/16 inch wide. Soregaroli (1971) has considered the quartz-potash-feldspar veins to be older than biotite veins. The writer, however, has consistently observed quartz-potash-feldspar veins transecting sulfide bearing biotite veins. Equally consistently the writer has found a second class of biotite veins identical to the above mentioned biotite veins (excepting for a lack of visible sulfide content) that cut quartz-potash-feldspar veins as well as quartz-molybdenite veins and some intramineral dikes. The later biotite veins are grouped in the fourth stage of mineralization and are discussed separately in a later section of this chapter. The biotite is dark brown or black with a felted texture and normally covering the fracture walls completely. Chlorite is generally present, the amount depending on the extent of alteration. Chalcopyrite and minor amounts of pyrite are the dominant and almost exclusive ore minerals; molybdenite and bornite, occur much less abundantly. All biotite veins combined probably account for less than 5 percent of the ore due mostly to their small size and low sulfide content.

### Mineralogy

Chalcopyrite appears as patchy, scattered blebs on the biotite walls, normally not covering more than 10 percent for the wall area. Aggregates of grains are generally less than 7 mm wide. Individual grains are as small as 0.1 mm.

Pyrite, the most abundant associated sulfide, appears to have




Figure 12. Contoured stereonet and major planes of orientation of veins.

been deposited either contemporaneously with or slightly later than chalcopyrite. Although grains or crystals are normally subhedral to anhedral, euhedral crystals are not uncommon. Most cubes and anhedral grains are 0.5 to 1 mm in diameter. Pyrite might form as much as 20 percent of the total sulfides, but normally is about 10 percent or less of total sulfides in Stage I veins. Although the percentage of pyrite versus chalcopyrite appears high in this system, it does not appear to seriously affect mill recovery because the veins are not important contributors to ore.

In the field bornite was identified in a single instance in a biotite vein by the writer but has not been observed in polished sections.

Macroscopic molybdenite is present in only a few biotite veins. Where it was observed, it is scattered as individual small plates or small groups of plates. Amounts are similar to the pyrite and in places equal to chalcopyrite. Although chalcopyrite-pyrite veins are earlier than second stage veins, most of the fractures bearing molybdenite were crosscutting and apparently later. It is suggested that these few veins, less than one percent of all biotite veins, are transitional to, or overlap in time the introduction of the second stage of minerals. Neither molybdenite nor bornite appear on a specific set of veins or in any particular orientation.

#### Structure

Although they represent 24 percent of all the ore veins measured (Stage I, II, and III), Stage I veins probably account for no more than five percent of the total sulfide ore due to the



minor concentrations of chalcopyrite and almost total lack of molybdenite in the veins.

No offsets were observed where biotite veins intersected aplite or pegmatite veins. Lineations, i.e. streaking of the felted biotite, are minor but present on some of the larger veinlets. Most lineations, however, are felt to be very late, associated with uplift. Lineations are commonly vertically oriented showing normal dip-slip movement when observable. Some also displayed horizontal strike-slip motions with both right-lateral and left-lateral movement. Lineations are also more abundant near shear zones though not tremendously so.

There are two concentrations of biotite vein attitudes. By far the major concentration strikes about  $N62^{\circ}E$  and dips  $78^{\circ}SE \pm 2^{\circ}$  (Figure 12a). It is outlined by two 15 percent concentrations out of 100 veins measured over the entire pit. The second high has an attitude of  $N76^{\circ}W/dip 86^{\circ}SW$  and was defined by visually centering the ten percent contour level. Orientation of veins are not found to change significantly from location to location (Figure 13). Preferred orientation for mineralogically distinct veins, i.e. molybdenite or bornite bearing veins, does not exist as far as present data allows. Soregaroli (1971) has described two orientations common to biotite-chalcopyrite veins: "a)  $5-20^{\circ}W/45^{\circ}E$  to vertical" and "b)  $N40-60^{\circ}W/15-30^{\circ}N$ " (Soregaroli, 1971). Because the writer has subsequently divided biotite veins between two stages (I and IV), the patterns obtained are expectedly different. A minor occurrence of veins striking  $N25^{\circ}E$  does exist, however, in the northeast section of the pit (Figure 13). Fracture densities,

Figure 13 Distribution of Stage I vein orientations within the Brenda pit. Strike and dip of major concentrations are heavy lines, and minor (questionable) concentrations are light lines. (Schmidt net used for construction)

Primary orientation is  $N62^{\circ}E$ , dip  $78^{\circ}SE$ . N indicates the number of observations in each sector. Dotted line indicates approximate pit outline as of August, 1971.





Figure 18 Distribution of Stage II vein orientations within the Brenda pit. Strike and dip of major concentrations are heavy lines, and minor (questionable) concentrations are light lines. (Schmidt net used for construction)

Primary orientation is  $N66^{\circ}E/78^{\circ}SW$ . N indicates the number of observations in each sector. Dotted line indicates approximate pit outline as of August, 1971.



measured in terms of number of fractures per yard, are commonly high. Due to their extremely small size, many density counts were only best guesses based perhaps on a few inches or a foot of exposed rock. Even with good exposure, many were not visible unless accidentally broken along their plane. Blasting dust also obscured veins making accurate counts and measurements difficult. Due to this difficulty, the writer cannot be certain if Stage I veins are more abundant in one locality than another in the mine.

### STAGE II VEINS

Quartz-potash-feldspar veins are distinctly later than biotite veins and constitute a second separate phase of sulfide mineralization. They are the most important source of ore, probably accounting for close to 80 percent of all sulfides. The reasons are: larger size veins, greater amounts of chalcopyrite and molybdenite in individual veins, and greater abundance of the veins, especially near the core of the deposit.

Vein thicknesses range from 1/8 inch to rarely 1&1/2 inches wide. The average size is about 3/16 of an inch. Generally sulfides comprise 10 to 30 percent of the vein space, but locally reach 100 percent.

### Mineralogy

Abundant hypogene sulfides are chalcopyrite and molybdenite. Magnetite, hematite, and bornite appear less commonly as hypogene minerals in Stage II veins. Most veins are closed. Open vuggy veins occur more commonly near the higher grade mineralized zones but possess exactly the same mineralogy and vein characteristics.

The bulk of the veins are filled with quartz unless sulfides are abnormally abundant. Early quartz is invariable anhedral forming grains as large as 3/8 inch. Rarely quartz has formed doubly terminated crystals exceeding an inch in length and embedded in chalcopyrite.

Potash-feldspar forms euhedral to subhedral crystals up to 1 cm wide projecting into the quartz along the vein walls. It also forms as alteration of wall rock that will be considered in more detail in a subsequent chapter. Disseminated chalcopyrite and pyrite occur rarely where the potassium alteration is particularly intense.

Small euhedral crystal aggregates of clinozoisite (identified by x-ray) are found sporadically upon quartz or chalcopyrite. They rarely have lengths of one inch, although most are much less than a centimeter long, commonly with a very fine-grained, anhedral layering. They are for the most part very late in paragenetic sequence.

Molybdenite is generally the first mineral deposited, either along the walls or as scattered, small blades in the quartz. Locally small stringers of molybdenite cut chalcopyrite. Grain size depends to a large extent on the total amount of molybdenite in the vein. Most blades are less than 1 mm to 1 cm wide. Molybdenite might be found in contact with any of the other minerals; and contrary to previous thought, much of it contains intimately, admixed chalcopyrite, and less commonly pyrite and bornite. Disseminated molybdenite is uncommon but is found rarely in intense hydrothermal biotite halos adjacent to mineralized veins (Figure 29).

Chalcopyrite is equal to or in greater proportion than molybdenite. Normally it is paragenetically later than molybdenite, forming in the center of veins. Blebs of chalcopyrite occur scattered over the vein walls, not interconnecting unless unusually high grades of mineralization are present (Figure 15).

Where chalcopyrite and molybdenite are in contact, the molybdenite is replaced in part along (cleavages) planes or between individual plates. These small wedges or laths are intimately mixed with molybdenite and commonly are as small as one micron (Figure 24). Pyrite is in places corroded by chalcopyrite. Fractures within pyrite grains are generally filled with chalcopyrite. Small laths and wedges of pyrite in molybdenite also occur but with less frequency.

Other veins show opposite relations. Pyrite in some veins forms euhedral cubes and pyritohedrons in chalcopyrite, which are often found in associations with euhedral epidote and doubly terminated quartz crystals, all embedded in masses of chalcopyrite (Figure 19). No specific orientations for this latter association of euhedral crystals was found. However, all were found close to high grade mineralization.

Bornite and sphalerite are less abundant constituents. Bornite is seen only rarely in hand specimens, but mostly occurs as very minute replacements along fractures in and along grain boundaries of chalcopyrite. It is generally less than 1/10 to 1/50 mm. Sphalerite has been seen only in polished sections in association with chalcopyrite. It occurs as rounded blebs or stringers up to one mm in diameter on the margins or in the

interior of chalcopyrite. Extremely minute disseminated grains of chalcopyrite within sphalerite give the appearance of exsolution textures (Figure 22).

Pyrrhotite has been reported previously to occur in association with chalcopyrite and pyrite; presumably in Stage II veins (Vogan, 1966). The writer, however, has studied many polished sections and has failed to observe any pyrrhotite. As far as the writer knows, pyrrhotite has not been reported since 1966; and then it possibly was misidentified.

### Structure

The predominant strike and dip of Stage II veins (taken from stereogram plots of poles to planes for 251 attitudes) is  $N66^{\circ}E/78^{\circ}SW$ , and remains relatively persistent throughout the mine (Figures 12b and 18). An exception is the young vuggy veins which strike mostly  $N0^{\circ}$  to  $10^{\circ}W$  and dip vertically. These are few in number and occur mostly towards the center of the orebody. They have also been observed by Soregaroli (1971) who noted a second orientation about  $N30^{\circ}-45^{\circ}E/15^{\circ}-25^{\circ}NW$ . The writer has not found a significant concentration in this range for the vuggy Stage II veins, however.

Densities of veining range from 5 to 15 fractures per yard along the periphery to about 70 per yard and occasionally well over a hundred per yard near the center of the orebody. A definite zoning of the fracture density of quartz-potassium-feldspar veins is the main control of ore grade. This aspect is discussed further in the following chapter.

Minor offsets occur here and there, but most were observed in blast debris; hence, true offset is mostly indeterminable.

Because of the steep dips of all fractures, it is most probable that movements were strike-slip since a large amount of movement is required for the same throw if dip-slip movement occurred rather than strike-slip. Maximum offset is normally less than one inch.

### STAGE III VEINS

The youngest ore veins at the mine are large quartz veins. Although not nearly abundant as earlier veins, they enhance the higher grade ore due to their larger widths and greater sulfide content. Size, properties of the quartz, and lack of potassium feldspar typically characterize and distinguish Stage III veins. Vein width averages one to two inches, although they can be as small as 1/2 an inch or rarely as large as 14 inches.

### Mineralogy

Hypogene ore minerals include molybdenite, chalcopyrite, pyrite, magnetite and sphalerite. Bornite has not been observed in these veins to date. Gangue minerals are normally quartz and minor amounts of calcite. Hematite occurs in small amounts associated with calcite. Any of the sulfides can occur singly or in combination with others and in widely variable proportions. Banding of the quartz and sulfide(s) might or might not be present.

Banded quartz-molybdenite veins consist of 1/4 to 3/4 inch repetitious layers of quartz separated by thin lacy molybdenite covers (Figure 16). The quartz is generally greyish due to disseminated molybdenite, though this is less common. Books of



subparallel molybdenite are as large as 1/8 inch, but normally do not exceed 1/16 of an inch in greatest dimension. Chalcopyrite and pyrite commonly are minor constituents forming irregular blebs of various sizes and tiny, locally-crosscutting veinlets, indicating they are probably paragenetically later than molybdenite. Pyrite forms both as striated euhedral cubes and anhedral grains. Cube edges commonly are less than 1/8 inch but can be as large as 1/4 inch.

Less commonly, the banded vein consists of chalcopyrite with molybdenite confined to the very earliest outer margin of the vein. Although there is a change in the predominant sulfide, the shape and structure is similar to the banded quartz-molybdenite veins.

A second type of Stage III veins is massive quartz-chalcopyrite or quartz-pyrite vein. The writer does not intend that a genetic difference be implied in this distinction - only a textural and mineralogical difference. All other features dictate the same system of mineralization. Chalcopyrite and pyrite occur as massive lodes in the vein or as smaller blebs scattered among the quartz. Molybdenite is normally missing or in subordinate amounts confined to the walls of the vein. Either chalcopyrite or pyrite can be the most abundant sulfide. Pyrite forms many euhedral, striated cubes with edges up to one inch long. Where present with chalcopyrite, most pyrite is situated along the margins of the vein. Locally, magnetite forms large veins with minor amounts of pyrite up to two inches wide. Euhedral magnetite grains massively packed together have maximum widths of 1/8 inch. Some of these veins have a larger amount of sphalerite (in the chalcopyrite)

than previous stages. These veins are more rare and are grouped with Stage III veins due to their size and similarity of occurrence, in the host rock. Most magnetite, where present, is less than one or two percent of the metallic content of the veins (Figure 21). Several samples of whitish "pyrite" were x-rayed to test the possibility they might be marcasite; all proved to be pyrite.

### Structure

The predominant vein attitude is  $N67^{\circ}E/77^{\circ}SE$  - defined by a 20 percent concentration of poles to planes. A secondary concentration occurs at  $N76^{\circ}W/86^{\circ}SW$  (Figure 12c).

Stage III veins are never densely spaced unless directly adjacent to a shear zone. In the core of the pit, densities reach one to two per yard though not consistently. Shears have been observed to transect these quartz veins but also may be unaffected by some. Offsets resulting from the development of the shears have not been observed by the writer. Similarly, offsets have not been observed due to the formation of the quartz veins. However, in a few places lensing of the quartz veins forms a sigmoidal shape. The similarity in orientation between vein stages suggests a similar structural origin.

#### STAGE IV VEINS

The fourth stage of mineralization consists of numerous veins and fractures that fall into four convenient types; clinozoisite veins, biotite veins, quartz veins, and calcite veins. All except a few of the early clinozoisite veins are post ore and intersect all previously described veins and intramineral dikes. It is possible that a hiatus is present sometime after the ore veins and before the Stage IV veins except epidote. Biotite veins are the most numerous and were at first mistaken for Stage I biotite veins as discussed earlier. They are quite thin, often discontinuous, not persistent, and difficult to observe unless the rock was fortuitously broken along one of the fractures. Minor amounts of sulfides are present especially where the veins intersect mafic dikes. Chalcopyrite, pyrite and hematite have been observed in Stage IV veins in generally subordinate amounts. Minor quantities of molybdenite have been observed in a few veins - especially clinozoisite veins.

Epidote veins (confirmed by x-ray to be clinozoisite) are for the most part insignificant in both number and sulfide content. Vein widths range from as small as 1/32 of an inch to 4 inches. Normally they are present as irregular 1/8 inch veins of finely granular anhedral clinozoisite (Figure 17). In the larger veins chalcopyrite, molybdenite, magnetite, (+ hematite), and pyrite might be present in greatly varying amounts and grain sizes. The most common assemblages of the larger veins are chalcopyrite-pyrite, chalcopyrite-molybdenite-pyrite, magnetite-pyrite, and magnetite alone. All except chalcopyrite occur either as anhedral

or euhedral crystals and crystal aggregates. Small anhedral grains of quartz can be interspersed among the epidote but never in amounts greater than 10 percent of the vein. Paragenesis of the sulfides is indeterminable by the writer, but it is likely that overall mineralization follows the pattern established for the preceding two stages.

Only 23 epidote veins were measured in the pit, insufficient data to give any concentration when plotted on a stereogram. Offset along these veins has not been observed. Soregaroli (1971) has observed "Epidote-magnetite-molybdenite veins which strike N45-55°W with vertical dips".

Clinozoisite veins are grouped in the fourth stage of mineralization but are in part contemporaneous with earlier veins. The larger veins are thought to be concurrent with the last phases of Stage III mineralization because the sulfide assemblages are similar, and the rare large lenses of sulfides are associated in space with some late Stage III quartz veins. The larger veins also show strong hydrothermal biotite envelopes very similar to Stage III alteration. Field observation has shown that the smaller veins and veinlets are younger than mafic dikes and hence are post ore. A possible mineralization hiatus might be present between the third and fourth stages of mineralization, although other structural features tie them together.

Biotite veins similar to Stage I biotite veins are approximately 1/32 inch to 1/16 inch but do not contain sulfides. For the most part, they are merely coatings of a dark brown felted mass of hydrothermal biotite on fracture surfaces. They are by far the most numerous veins of this stage. Rarely pyrite and/or chalcopyrite

occurs as isolated irregular blebs less than 2 mm long.

Small quartz veins, 1/4 inch or less in width, crosscut all previously mentioned mineralization. These are few in number and contain no related potash-feldspar, biotite or other alteration minerals. In places they contain euhedral pyrite or minor amounts of anhedral chalcopyrite. Sulfides are rare and were observed only where crossing post-ore dikes. They do not appear in any way related to ore veins. In one case a vuggy, though unmineralized quartz vein, was observed crosscutting the trachyte dike.

Calcite veins are numerous though small in size. They fill in most reopened fractures of all ages and form their own veins, commonly 1/8 inch wide and rarely two to three inches. Hematite is ordinarily the only opaque mineral in these veins. It is thought to be disseminated among the calcite, producing a black-brown to orange tinge. Positive identification of the calcite coloration was not possible due to the extremely fine-grained and dispersed nature. Rarely crystalline ferroan dolomite has been found to fill open fractures at a very late stage.

Specular hematite has, at rare localities, filled small 1/16 inch fractures. Their exact relation to other veins is unknown.

A single barite vein, one to one and one-half inches wide, occurs in the northwest margin of the pit. The margins of the veins were brecciated with quartz diorite fragments from 1/2 inch to one inch in size. This is the only observed occurrence of barite in the pit. It is in an area of hydrothermal biotite alteration, but the vein itself was found loose in the muck; and the writer was unable to locate its outcrop.

### Structure

Fourth stage veins (dominantly biotite veins) are similar in attitude to the previous veins. High concentrations on the Schmidt net are N66°E/80°SE and N64°W/89°SW (Figure 12d). Greater scatter of attitudes exists among these veins than in other vein systems; possibly because they are late and formed in a more complicated environment than ore fractures. But a consistent change in attitudes relative to location in the pit cannot be determined. The fracture density of the biotite veins in places appears high, but a precise count was not always possible. Calcite and quartz veins are most numerous around dikes and peripheral to shear zones.

### SECONDARY MINERALS

Secondary minerals observed include ferrimolybdate, limonite, malachite, azurite, chalcocite, native copper, powellite, cuprite, cupriferous manganese oxides, and ilsemanite. Tenorite and covellite have also been reported by earlier workers (Soregaroli, 1971).

Limonite is the predominant secondary mineral, forming in weathered rock that extend rarely to depths of 90 feet below the surface. The average weathered zone where limonite forms is less than 20 feet thick. Greater depths of weathering and hence of limonite formation are found locally in shear zones. Limonite forms as thin, brown coats and in some cases as boxwork replacement of pyrite and chalcopyrite. X-ray analyses did not reveal any goethite.

Ferrimolybdate, an alteration product of molybdenite, is uncommon at Brenda and has been identified by the writer in only two samples. Samples were, however, more easily observed when early

stripping of the cap rock was taking place (Soregaroli, personal communication). One sample was found in a one inch quartz-chalcopyrite-pyrite vein located a mile northeast of the mine. The other was found at the tailings dam and was presumed to have been hauled from surface muck obtained at the pit. Favorable conditions for the formation of ferrimolybdate require a low pH on the order of 1.5 to 4. The absence of significant amounts of pyrite and a cold climate both could be responsible for the overall scarcity of this alteration product.

One small specimen of native copper was observed in a late vuggy calcite vein.

Malachite and azurite have been found as thin encrustations along joints or fractures in the uppermost exposed section near the surface. In places the encrustations are 1/8 inch thick, but normally they are much thinner. Malachite is more abundant than azurite.

Negligible supergene enrichment is recorded in Brenda deposit. Admittedly, however, it is only a single occurrence. In a single location, sample #4-3 of bench 5060 located near the surface of the old stream which transversed through the center of the pit, sooty chalcocite was confirmed as a thin coating on pyrite and chalcopyrite grains along a Stage II vein. A layer of chalcocite one to three mm thick in places almost completely replaces 1/2 cm chalcopyrite and pyrite grains. The extent of this supergene enrichment is limited to only a few yards near the former streambed where this particular vein is exposed. Pyrite and magnetite, it should be noted, form as much as 20 percent of this vein; and




Figure 25. Line diagram of mineralogy of vein systems. The thickness of the lines indicates general relative abundance of the minerals.



pyrite is in great abundance in an 8 inch quartz-pyrite vein immediately adjacent. Hence, many of the conditions appear to have been combined fortuitously for this rare supergene occurrence.

#### HYDROTHERMAL ALTERATION

Hydrothermal alteration in the Brenga ore body has been described in more or less general terms by several previous workers (Vogan, 1966; Carr, 1967; and Soregaroli, 1968 and 1971). While differing significantly from the first two authors, the writer's alteration study is in agreement with the types of alteration presented by Soregaroli (1968 and 1971). These types are 1) potassium alteration, 2) propylitic alteration, and 3) argillic alteration. The first two types form as narrow envelopes about Stage I and II veins, while argillic alteration is associated with Stage III veins in zones up to 35 feet wide and several hundreds of feet long. Although searched for dilligently, a spatial pattern of alteration, other than the small envelopes about ore veins, could not be found. Concentric zoning of alteration types, for example, does not exist as far as the writer has been able to determine. Soregaroli (1971) indicates the presence of hydrothermal biotite in a zone extending to the northeast beyond the ore body.

Potassium alteration consists of two types: potassium feldspar (chiefly orthoclase) and hydrothermal biotite. Potassium feldspar mostly forms thin envelopes around Stage II veins (Figures 26, 27, 28, and 31). Generally the feldspar is confined to growth along vein walls but can form envelopes in the wall rock up to 8 inches wide replacing plagioclase in the host rock. Occasional mica and kaolinite have been seen replacing mafics and some feldspar.

Where large amounts of potassium feldspar are introduced, the hydrothermal biotite content also increases and forms plates up to 1/4 inch diameter admixed with the feldspar. A secondary envelope of hydrothermal biotite might also develop around the feldspar envelope before grading into propylitic alteration. Although hydrothermal potassium feldspar is associated with the most important ore veins (Stage II), the amount of potassium feldspar does not appear to be related to the amount of sulfides present.

Hydrothermal biotite, often a green biotite under the microscope, occurs in all stages of vein formation. Only in the second stage is it less developed. In the first stage of mineralization, it forms felted black coatings as described earlier. Rarely is it visibly developed into the wall rock adjacent to the vein. In second stage veins it is less developed but can form a thin envelope; generally less than 1/8 inch and difficult to observe. Macroscopically, it can be recognized by its greater abundance and its fine-grained nature compared to normal host rock biotite. It also may be admixed with orthoclase as mentioned earlier or form a secondary envelope about the potassium feldspar alteration (Figure 31).

Third stage veins also contain abundant biotite and generally little or no potassium feldspar. Commonly, alteration is so complete that hydrothermal biotite forms more than 50 percent of the rock. This envelope can be several feet wide in some places, while elsewhere it is less than one inch wide. Rarely disseminated sulfides are present in the biotite envelope (Figure 29).

Stage IV alteration consists of biotite filled fractures similar to Stage I veins and envelopes of biotite a few inches wide around

large epidote veins which are generally few in number. In all cases biotite and potassium feldspar grade outward into propylitic alteration.

Propylitic alteration is a term used to describe a weak alteration effect forming thin envelopes adjacent to potassium alteration along veins. Where vein densities are extremely high, pervasive propylitic alteration might be present, though not necessarily. Epidote and minor amounts of chlorite are characteristic alteration products of the propylitic zone. Epidote commonly forms in the cores of plagioclase crystals, and occasionally muscovite appears in the assemblage. Megascopically, the plagioclase appears slightly greenish and waxy textured.

Thin white albitized veinlets occur rarely, but their exact relationship to other alteration types has not yet been determined. Indeed, they are possibly late weathering alteration unrelated to ore vein alteration.

Argillic alteration occurs in large widely spaced planar zones, generally a light tan or brown in color (Figure 8). Stage III veins are always associated, and shearing is generally present. X-ray analyses of clay fraction of nine alteration zones of various widths and locations show identical assemblages of kaolinite, muscovite and montmorillonite. For an assemblage such as this, Meyer and Hemley (1967) suggest using the term "intermediate" argillic alteration to distinguish it from a more advanced argillic alteration which they also describe.

Although both biotite and argillic alteration are associated with Stage III veins, the writer could not find a consistent spatial




Figure 32. X-ray diffraction traces clay fraction of argillic alteration zone. Kaolinite, muscovite and montmorillonite are dominant constituents. Numbers indicate degrees  $2\theta$ . KV = 40, MA = 20.

relationship between biotite and argillic alteration. It appears that only certain Stage III veins (generally two or three per shear zone) are suitable for producing this alteration. Most of the veins in these zones were high in quartz and molybdenum; however, some quartz-molybdenum veins were observed which were not associated with argillic alteration.

Muscovite, much of it fine-grained, occurs throughout the shear zone. In places it might be termed "sericite" alteration; however, because of its extremely rare and inconsistent appearances, the writer believes it does not warrant a separate classification.

#### SUMMARY

Ore minerals occur almost exclusively in veins at the Brenda ore deposit. These veins have a specific paragenetic order. Ore forming veins are biotite, quartz-potassium-feldspar-sulfide, and quartz-sulfide veins.

Biotite veins are the earliest. They are thin veins containing minor yet consistent amounts of chalcopyrite. Predominant orientation of these veins is N62°E/78°S and N76°W/86°S.

Quartz-potassium-feldspar veins formed later, apparently without a significant time gap separating them from the earlier biotite veins. Concentrated orientations occur at an attitude of N66°E/78°SE. Vein densities increase toward the center of the pit and are the principle "raison d'etre" for high grade ore zones. Average vein width is 3/16 of an inch. Strike-slip offsets are abundant. The ore mineral paragenesis is molybdenite, chalcopyrite, pyrite, magnetite and hematite, although some reversals do occur. Potassium feldspar is

the main alteration type usually grading into propylitic alteration. A thin envelope of hydrothermal biotite might also be present.

Stage III veins are large quartz veins containing various sulfides in several combinations. Banded molybdenite veins and massive chalcopyrite veins are the most common. Potassium feldspar is normally absent; however, biotite and argillic alteration are commonly associated. Primary orientations are  $N69^{\circ}E/77^{\circ}SE$  and  $N67^{\circ}W/86^{\circ}SW$ . They have in places the form of sigmoidal lenses.

Stage IV veins are post ore and contain no significant concentration of sulfides. They consist of clinzoisite veins, biotite-lined fractures, calcite and quartz veins. Large epidote (clinzoisite) veins overlap in time of formation with late Stage III veins. All other Stage IV veins and fractures are significantly later and probably distinctly separate from the third stage of mineralization. Attitudes are more widely scattered than in other stages of mineralization. A small 10 percent concentration occurs with an attitude of  $N66^{\circ}E/80^{\circ}SE$  and  $N64^{\circ}W/89^{\circ}SW$ . Offsets have not been observed by the writer.

All vein systems possess remarkably similar orientations that do not change significantly over the pit area. A slight progressive change in attitude is possibly indicated for Stage I to III veins inclusive; Stage I strike  $N62^{\circ}E$ ; Stage II -  $N66^{\circ}E$ ; Stage III -  $N69^{\circ}E$ . These small differences, however, might not be real and could result from the relatively small amount of data available. Stage IV veins are more scattered and have smaller concentrations, possibly due to later structural influence in the system long after the ore forming environment had passed.

Hydrothermal alteration is limited to envelopes about veins, except for a few large zones of argillic alteration associated with large Stage III veins. Hydrothermal biotite and potassium feldspar (chiefly orthoclase) form about Stage II veins, while biotite alteration forms in varying degrees about veins in Stages I, II and IV. Propylitic alteration might accompany all stages of veining in varying amounts, although generally it too is limited to envelopes about veins.

## CHAPTER V

## ZONING


MINERAL ZONING


At Brenda mine it is possible to find a general systematic distribution for abundances of chalcopyrite, molybdenite, and galena, although no such patterns have been recognized for other minerals such as bornite, sphalerite and pyrite. Nor have patterns been found for the abrupt appearance or disappearance of any of the ore minerals. Chalcopyrite and molybdenite zoning results from a change in the density of veins across the ore body.

Copper and Molybdenum

The presence of a central zone of high copper and molybdenum percentages has long been recognized. Because the only significant primary copper and molybdenum minerals are chalcopyrite and molybdenite, the grade distribution is directly related to mineral distribution. Figures 33 and 34 display the zoning pattern for copper and molybdenum grades respectively on the 5210 bench. Cuttings from closely spaced 58 foot deep rotary drill holes were uniformly sampled and analyzed by atomic absorption at the mine. Lines of equal abundance were hand contoured. A distinct, though irregular central zone of higher grades, exists in the center of the ore body. To the west the amount of copper drops quickly in an almost linear zone that coincides roughly with the appearance of large xenoliths of Nicola Group hornfels mapped by the writer. Reductions in copper and molybdenum take place more gradually to







the east. The limits of the present pit end in low grade ore on the eastern margin.

The increase of copper and molybdenum assays in the center of the ore body is caused mostly by a large increase in the density of Stage II veins accompanied by a slight increase in the average width of the veins. Stage II veins in the periphery of the pit average from 5 to 15 per yard; while near the center of the pit, vein densities may be as high as 100 per yard or more in some localities.

In addition, a notable increased variation of attitudes of Stage II veins also occurs in the central part of the deposit. Veins with attitudes between  $N35^{\circ}E$  to  $N40^{\circ}W$  occur more often in areas of higher total grade. The contribution of Stage I and III veins appears to be relatively less important to the zoning in the pit. Indeed Stage I veins seem to decrease in number in the center of the deposit relative to the periphery.

#### Lead

To date only a small part of the 5110 bench drill holes has been assayed for lead. Most lead in the mine is less than 0.005 percent, but small isolated patches up to 0.2 percent appear here and there.

Analyses for lead on the 5060 and 5110 bench were performed on drill cuttings in a similar fashion to copper and molybdenum analyses. A statistical treatment of 323 values was done in order to separate possibly distinct populations according to the methods of Tennant and White (1959), Lepeltier (1969), and Sinclair (1972). The cumulative percent frequencies were plotted on cumulative log

probability. Examination of the resulting curve suggests that two lognormal populations are represented in the data. Population R forms 90 percent of the total, and Population S forms 10 percent (Figure 35). Recalculation of the values to 100 percent (for each population where not overlapping) defines the distinct end member population (Symbol ● in Figure 35). Because these are two interpreted lognormal populations, it is a useful check to mix the two in appropriate proportions for comparison with the real data. When mixed in the ratio 10 to 90 (for S and R respectively), the resulting curve (Symbol o in Figure 35) is seen to match closely the real data (Symbol X in Figure 35).

The R population, with a geometric mean of 32 ppm, can be interpreted as rock background values contained in K atomic sites in such minerals as potash-feldspar, biotite, and hornblende. The value compares with a world average of 15 ppm in intermediate intrusive rock (Krauskopf, 1967).

The S population probably represents sporadic localities of sulfide lead, for minor amounts of galena have been seen in the mine.

The distribution of these values is shown in Figure 36. The S or assumed sulfide population occurs as isolated sporadic highs. Since only a limited area of the pit has yet been sampled, interpretation of a pattern is highly uncertain. Possibly fracture density might also relate directly to sulfide lead; and if so, an expected high in the center of the pit should exist. A second possibility is that galena increases away from the center of the deposit; however, present data is too limited to confirm or deny




Figure 35. Cumulative probability plot (M) of 323 blast-hole analyses of lead (ppm) from Brenda Mine. Lines R & S are the populations separated from M. Open circles represent ideal mixing points of R & S in the proportions 90% R and 10% S.




Figure 36. Distribution of Lead in Brenda pit from blast-hole analyses.

this possibility. Additional analyses are underway at the mine to further examine the distribution of lead in the mine area. Other minerals common in and about the ore zone are bornite, magnetite, hematite and pyrite. Of these only pyrite appears to show even a vague systematic variation in abundance, apparently increasing in amount to the north and west.

#### MINOR ELEMENT ZONING

A study of minor element distribution in 50 chalcopyrite samples from Brenda mine has been undertaken by Dr. D. Brabec. Samples, provided by the writer, represented the veins of Stages II and III. Samples were analyzed for Se, Ni, Cd, Mn, Zn, and Ag. The latter five were analyzed by atomic absorption spectroscopy, whereas selenium was analyzed by x-ray fluorescence. Geometric means for these elements (lognormal distribution) were Cd - 31ppm; Mn - 87ppm; Ni - 16ppm; Ag - 79 ppm; Zn - 734ppm; and Se - 358ppm. Plotted results do not indicate the presence of lateral systematic variations (i.e. zoning). However, sample distribution was erratic; and the number of samples was relatively small compared to the size of the pit. Hence, a possibility of other perhaps complicated patterns cannot be ruled out. Significant positive correlations at a probability level of 0.05 were found with Cd and vein width; Cd and Cu/Fe ratio and Ni and Ag. Negative correlations were found for Ni and Cu/Fe ratio; Zn and Se; and Se and vein width. Although not all have been explained, cadmium apparently increases in thicker veins of Stage III type, whereas selenium decreases in these later veins compared to Stage II veins. Except for Cd and Ag, the elements

show a bimodal distribution on lognormal probability plots

(D. Brabec, personal communication, 1972).



## CHAPTER VI

## SUMMARY AND CONCLUSIONS

GENERAL GEOLOGICAL EVOLUTION OF THE BRENDA DEPOSIT

Several textural varieties of quartz diorites form the Brenda Stock that was emplaced in Nicola Group volcanic rocks about 176 m.y. ago. One phase, the Speckled quartz diorite, contains most of the Brenda copper-molybdenum ore body. Small aplite and pegmatite dikes formed shortly thereafter, probably during a late stage of the cooling history of the intrusion.

Contrary to Carr (1967), the writer found that these felsic dikes, although more abundant near the central part of the pit, formed in all areas of the ore body and are not confined to a structural belt as proposed by Carr (1967). Furthermore, these orientations do not correspond to ore veins or other younger planar features. Consequently, the writer is inclined to relate these felsic dikes genetically to the Brenda Stock as a late stage residuum. This dissimilarity between felsic dikes and all younger features also supports the hypothesis (contrary to Carr, 1967) that a period of time, probably large, elapsed after the intrusion of aplite-pegmatite dikes and before fracturing and mineralization of the ore body began. Potassium-argon age dating data preferentially support (in the writer's view) the hypothesis that mineralization occurred at about 146 m.y. - well after the solidification of the intrusive quartz diorite at about 176 m.y. On this assumption it appears highly unlikely that this deposit is a paramagmatic type deposit,

i.e., "an integral feature of a magmatic event." (White et. al., p. 9, 1968).

The Brenda ore body was formed by three separate but partially overlapping pulsating phases of mineralization. An increase in fracture density in the central part of the pit causes an irregular zoning of copper and molybdenum grades at Brenda. There is, however, an absence of symmetrical lateral zoning of minerals and elements in the classical sense and a scarcity of hydrothermal alteration, other than as thin envelopes about veins. Because vein orientations remain rather uniform throughout the deposit and because of their strongly preferred orientation and similarity to air photo lineament directions (Chapman, 1968), the writer believes that a regional rather than a local control caused fracturing that provided channels for migrating ore fluids.

Although some movement along fractures occurred during mineralization as evidenced by minor offsets of veins, most movement began near the end of the third stage of mineralization. Shearing took place in weakened zones of argillic alteration and faulting, both dip-slip and strike-slip followed. The first minor epidote veining also took place at the end of the third stage of mineralization, but most Stage IV veins developed after the emplacement of the intramineral dikes. Whole rock analyses of two intramineral dikes indicate an age of about 130 m.y.

Late Tertiary events are largely speculative. Movement along shear zones and perhaps some faults probably continued during the Tertiary but did little to affect the ore body. A small patch of Tertiary (?) conglomerate possibly once extended over a large area

but has been largely eroded. Glaciers might have eroded appreciable portions of the ore body. While at the surface and subjected to weathering and weathering processes, produced features such as the black clays developed from the combined breakdown of the argillized zones and the physical transport of fine clays, molybdenite and other opaques downward along accessible fractures. Ferrimolybdite, limonite, malachite, powellite, and other secondary minerals developed in a zone of weathering that averaged 20 feet deep, but locally attained depths of 90 feet.

#### UNUSUAL ASPECTS OF THE BRENDA DEPOSIT

The presence or absence of a number of features sets Brenda apart from other large tonnage, low grade copper and/or molybdenum deposits.

1) A well-defined spatial alteration and ore mineral sequence is virtually absent. Potassic and propylitic alteration is limited to thin envelopes around vein walls. Even the highest grade ore can have practically no visible alteration affects, whereas some almost barren veins have prominent envelopes up to six or eight inches wide. The greatest alteration effect, argillic alteration, occurs in widely spaced planar zones no wider than 35 feet and generally associated with a few third stage veins and shearing. What a contrast this is to the alteration at Climax or Bingham or most large tonnage deposits one can name. That the deposit also appears to lack a classical zonation of ore minerals (other than that related to density of fractures) compounds the disparity.

2) Unlike most similar deposits, disseminated ore mineral occurrences are extremely limited at Brenda. All occurrences of disseminated sulfides observed by the writer occur in zones of unusually intense potassic alteration, either hydrothermal biotite or potassium feldspar. It is probable that the lack of disseminated mineralization is related to the scarcity of alteration particularly potassic alteration.

3) Another unusual aspect of the deposit is the remarkable uniformity of orientation of structures with time. From the first phase of mineralization through shearing, faulting, and diking to the last phase of mineralization only two orientations persist. The dominant orientation strikes N60-70°E, and the second orientation strikes N55-65°W. Both have steep southerly dips.

4) Also differing from most "Porphyry Copper" or "Stockwork-moly" deposits is the approximately equal economic grades of copper and molybdenum at Brenda. Most deposits of this type predominate in either copper or molybdenum but rarely both. Rhenium values apparently support this "middle of the road" status of Brenda. The rhenium content of molybdenite concentrates are about 120 to 180 ppm at Brenda, whereas most porphyry copper deposits have high values (Ely - 1650 ppm, Chino - 800 ppm, Bingham Canyon - 360 ppm); and stockwork molybdenum deposits are generally quite low in rhenium content (Questa - 12 ppm, Climax - 3 ppm), (Sutulov, 1970).

## PROBLEMS FOR FUTURE INVESTIGATION

A particularly important problem revealed by this thesis is the need for additional potassium-argon analyses for final determination of the ages of the quartz diorite host and ore mineralization. Present data indicates that mineralization occurred at a maximum of 146 m.y. and possibly as late as 130 m.y. It is critical to analyze biotite strictly related to ore mineralization (Stages I, II, or III) being careful not to include Stage IV biotite alteration that already is indicated to be 130 m.y. In addition, further samples from entirely unmineralized areas and from selected intramineral dikes are needed for confirmation.

Careful regional mapping is needed to tie the structural and petrographic history into a regional picture. Too little is known about the Brenda Stock. This vicinity has been termed by some to be part of the Pennask batholith (Schau, 1970) and by others to be part of the Similkameen batholith (Peto, 1970). Internal details such as phase relations and contacts need careful mapping, for the writer has found that the relations between varieties of quartz diorite might be much more complex than previously mapped. Careful attention should also be paid to secondary structural features such as veins, dikes faults, and shears in order to try to relate them to features observed within the Brenda ore body. In addition, the Nicola Group rocks are little understood in the vicinity of Brenda and deserve consideration.

A project for future consideration is an element zoning study from blast-hole data. Brenda is presently beginning to store copper and molybdenum concentrate from each blast hole (D. F. Whitford, Brenda Mines, personal communication). Considering the tight

patterns of blast holes, it might be possible at a later date to study both lateral and vertical zoning with thousands of samples available for analyses by x-ray fluorescence.

It has been suggested by Dr. Soregaroli (personal communication) that because of the limited alteration present at Brenna mine, it might be much more amenable for detailed study than deposits where alteration is pervasive and overlapping. The writer believes that a detailed study of the nature, chemistry, petrology and interdependence of the wall rock alteration assemblages would be a valuable contribution.

## BIBLIOGRAPHY

- Carr, J.M. 1967. Geology of the Brenda Lake Area. B.C. Bureau Mines. Mines and Pet. Res. Report. pp. 183-210.
- Chapman, Wood and Griswold, Ltd. 1966-1967. Brenda Feasibility Study. Brenda Mines Ltd.
- Chapman, E.P. 1968. Geology of the Brenda Molybdenum and Copper Deposit. Canadian Inst. Min. Metal. Bull. April, 1968. Vancouver, B.C.
- Drummond, A.W. and Kinura, E.T. 1969. Hydrothermal Alteration at Endako - A Comparison to Experimental Studies. Canadian Inst. Min. Metall. Bull. Vol. 62, p. 707-714.
- Fountain, D.K. 1968. The Application of the Induced Polarization Method at Brenda Mine, B.C. Inst. Min. Metal. Bull. Vol. 61, No. 670, pp. 153-157.
- Hawley, J.E. and Nichol, I. 1959. Selenium in Some Canadian Sulfides. Econ. Geology. Vol. 54, No. 4, pp. 608-628.
- Kirkham, R.V. 1971. Intramineral Intrusion and their Bearing on the Origin of Porphyry Copper and Molybdenum Deposits. Vol. 66. No. 8, pp. 244-249.
- Krauskopf, K.B. 1967. Introduction to Geochemistry. McGraw-hill. New York. p. 721.
- Lepeltier, C. 1969. A Simplified Statistical Treatment of Geochemical Data by Graphical Representation. Econ. Geology. Vol. 64, pp. 538-550.
- Little, H.S. 1961. Vettle River (West half). Geol. Survey Canada. Map 15.
- Menzies, M.M. 1969. The Brenda Project. Canadian Mining Journal. pp. 120-124.
- Neyer, C. and Hemley, J.J. 1967. Wall Rock Alteration in Barnes, h.L. (ed.), Geochemistry of hydrothermal Ore Deposits. New York. Holt, Rinehart, and Winston. pp. 166-235.
- Nasmith, H. 1962. Late Glacial History and Surficial Deposits of the Okanagan Valley, B.C. B. C. Dept. of Mines, Bull. 46.
- Ney, C.S. 1956. Unpublished company report. Northwestern Exploration, Ltd.

- Ney, C.S. 1957. Geological and Geophysical Report on the Brenda Prospect. Osoyoos Mining Division, B.C. Assessment Report No. 189.
- Peto, P. 1970. Unpublished M.Sc. Thesis, Univ. of Alberta.
- Rice, H.M.A. 1947. Princeton Map-area. Geol. Surv. Canada. Mem 243.
- Schau, M. 1970. Stratigraphy and Structure of the type area of the upper Triassic Nicola Group in South-Central British Columbia. Geol. Assoc. Canada. Special Paper No. 6. Structure of the Southern Canadian Cordillera. p. 123.
- Sinclair, A.J. 1972. Some Statistical Applications to Problems in Mineral Exploration. B. C. Professional Engineering. March, 1972.
- Soregaroli, A.E. 1968. Geology of the Brenda Copper Molybdenum Deposit, British Columbia. Paper presented at Northwest Mining Assoc., Spokane, Washington.
- Soregaroli, A.E. 1971. Geology of the Brenda Copper Molybdenum Deposit, British Columbia. Paper presented at Quebec City, IMM 73rd Annual General Meeting.
- Sutulov, A. 1970. Molybdenum and Rhenium recovery from porphyry coppers. Univ. Concepcion, Chile. p. 259.
- Tennant, C.B. and White, M.L. 1959. Study of the Distribution of some Geochemical data. Econ. Geology. Vol. 54, pp. 1281-1290.
- Vogan, R.S. 1966. Geological Report - Feasibility Period. Brenda Mines, Ltd.
- Wallace, S.R., Muncaster, N.K., Jonson, D.C., MacKencie, W.B., Brookstrom, A.A. and Surface, V.G. 1968. Multiple Intrusion and Mineralization at Climax, Colorado in Ridge, J.H. (ed.). Ore Deposits in the United States. Graton-Sales. New York. Amer. Inst. Min. Eng. Vol. 1, pp. 605-640.
- White, W.H., Harakal, J.E., and Carter, N.C. 1968. Potassium-Argon ages of some ore deposits in British Columbia. Canadian Inst. Min. Metall. Bull. Vol. 61, pp. 1326-1334.
- White, W.H., and Harakal, J.E. 1972. Personal Communication on unpublished Potassium Argon Age Data from the Brenda Mine Deposit.