

Energie, Mines et Ressources Canada Secteur de la Commission géologique du Canada

601 Booth Street Ottawa, Ontario K1A 0E8

25 June 1991

P. Lougheed Senior Geologist Prime Explorations 11th Floor, Box 10 808 West Hastings Street Vancouver, B.C. V6C 2X4

Dear Peter:

The Bondar-Clegg analyses for Tantalus samples, in general, especially as pertains to gold, are rather disappointing. Most samples are low in gold including the high pyrite, sartorite- and natroalunite-bearing samples (OHSA-545 and 546) and my high pyrite samples (KQ-90-180A, 180B, 182C and 182D) west of the Treaty Glacier.

The most encouraging samples are KQ-90-171A, 171B and 171C (spaced about 100m apart) and KQ-89-136B (about 200-300m north) near the Treaty Glacier on the western-most extremity of the Treaty Gossan (see attached location map). These samples of altered pyritic plagioclase-hornblende porphyry (diorite to monzodiorite?) cut by pyrite, quartz veinlets (impart leached-even in fresh outcrops adjacent ice) suggest a fairly large area anomalous in gold. This area is similar to parts of the Konkin Nunatak.

Samples KQ-89-122D and 122E (pyritic altered rhyolite-ridge crest northeast of main Treaty Gossan) and KQ-89-136B (pale quartz sericite, pyrite schist west side of main gossan, mentioned above) anomalous in gold are cut by 1 to 3% quartz veinlets. This relationship suggests to me that in prospecting the main Treaty Gossan that emphasis should be placed on finding and sampling veins.

We very much appreciate your and Prime's help in obtaining analyses on these samples. I just wish that we had been a bit more luckyin finding gold-rich samples! Perhaps this summer?

Sincerely,

R.V. Kirkham

RVK/lo

cc: J. Chapman

S.B. Ballantyne

