

## MOUNT ARMOUR

## \* I) GEOLOGIC STYLE - LITHOLOGY

- Map area dominated by major mafic volcanic sediment contact.

## i) Mafic Volcanics

- coarse to medium grained basalts, pillow basalts and possible sub-volcanic intrusions.

- hiatus in volcanism demonstrated by thin N/S trending chert unit bounded by mafic volcanics, western margin of property.

## ii) Altered Mafic Volcanics.

- Generally very light. Rare occurrences (cf. L35 900W) of light FeC development minor Py  $\pm$  sericite.

- Somewhat more widespread low grade cc development.

## iii) SEDIMENTS

- Generally dominated by a quartz rich regime

## @ Cherts + Ribbon cherts:

- Dominant over much of the ~~the~~ western portions of map area.

Bedding in RC variable From 5-100 cm.

- Well developed chert bx's are noted, these show evidence of healing

by secondary quartz but no sulfides. On no portion of the property was the equivalent of a Rea Bx noted.

- Between L45 - L55 275 W, the chert unit has been extensively altered by a Py - silica system. Suggestion has been forwarded that cherts in this region are acting as a cap rock.

- May appear anomalously thick due to folding eg. L 400 N - 400 W

Ⓒ QUARTZ - PEBBLE Congl.

- Appears to be the facies equivalent of the massive chert unit.
- Generally well sorted quartz and chert pebbles, rounded to sub-rounded, may contain (locally) slight disseminated hematite.
- No distinctive alteration.

Ⓓ QUARTZ - WACKES and QUARTZITES

- Widespread distribution, may contain occasional argillite interbeds. These rocks occur in close proximity to the main sulfide showing. Small slightly pyritized exposures at L3 + 05 S 75 W

Ⓔ ARGILLITES: (Phyllites + Mudstones)

- Widespread distribution possible association with main sulfide showing. Suggest, based on intuition (and without o/c evidence) that argillites may be thickening slightly in the vicinity of the main showing.

- Agillites, small o/c noted within 50 m of second (South) massive sulfide showing.

© LIMESTONE - LIMESTONE Cobble Bx

- DISTRIBUTION confined entirely to Northern portions of MAP AREA. Appears to form classical reef and fore-reef SYSTEM.
- Bx's may have LST Fragments in excess of 100 m.
- No significant alteration associated with this unit.
- Dirty LST lens infrequently noted within unit 63.

NOTE: All other map units are essentially combinations of the above.

II GEOLOGIC STYLE: STRUCTURE

- All structural data summarized on FIGURES II - IV.
- Two distinctive and related Fold styles
  - i) LARGE SCALE 90° WARP.
    - STRUCTURE best demonstrated by gradual changes in bedding attitudes.
    - Poles to bedding, Figure IV,

in combination with an estimated trend of the axial trace suggest the MAJOR FOLD is :

- 1.) SYMMETRIC ABOUT  $165^{\circ}$
- 2.) Upright EASTWARD FACING AXIAL Plane  $165/77E$
- 3.) Plunging NORTH AT MODGRATE  $35 - 45^{\circ}$  angles.

## ii) TIGHT Isoclines.

- A series of three <sup>(minimum)</sup> of these Folds are discernible in the N-S trending o/c ridge and cliffs  $L15 \rightarrow L55 \approx 300W$ .
- Most convincing structure at  $L1+305 - 325W$
- Significant that data of FIGURES II AND FIGURES IV, do not allow for the development of a 2<sup>nd</sup> Fold phase. These isoclines are compatible with the  $90^{\circ}$  WARP Fold, when viewed as smaller ancillary Folds OR DRAG FOLDS.
- Position of axial trace of these Folds is contingent on the relations, Facing, of unit 61 (chert) ; 67 (WACKES - Fine grained quartz wackes). Chert progressively overlies  $\rightarrow$  underlies  $\rightarrow$  (repeated) wacke unit, as cliff face is traversed.

III SULFIDE SHOWING - RELATION TO STRAT/STRUC.

- Strike relations derived from either pyrite fragmental "beds" (North Trench) or adjacent argillites (South Trench) and other <sup>spatially related</sup> attitudes, suggest that the two lenses represent stacked horizons, not a single folded horizon.
- Critical thickening of each of these pods may be occurring in the hinge region of the tight isoclinal, (35-40° to 335°).
- Some characteristics of stratigraphy analogous to Baschi type deposits. (Salton Sea analogue). May be interesting to check cobalt content of massive sulfide pods.

# FIGURE I

## GENERALIZED STRATIGRAPHIC COLUMN MOUNT ARMOUR.

The thickness of the lithologies in this column are calculated from average attitude and contact relations. Folds repetitions are not represented within this section.










FIGURE II

MOUNT ARMOUR LINEAR FABRIC



CONTOUR DENSITY  
N = 24

	5-10 %
	10-20 %
	20-30 %
	30-35 %
	35-40 %

CONCENTRATION OF  
LINEAR DATA AT :


42 → 336

17/06/84





50

→

FIGURE III  
MOUNT ARMOUR FOLIATION  
DATA.



CONTOUR DENSITY.

	2-5 %
	5-8 %
	8-10 %
	10-12 %

CONCENTRATION OF  
Planar SURFACES ABOUT  
Two POLES:

$$F_1 = 34 \rightarrow 203.5$$

$$F_2 = 12 \rightarrow 240.5$$


$$F_1 \quad 118/57 \text{ NE}$$

$$F_2 \quad 151/78 \text{ NE}$$







FIGURE IV

MOUNT ARMOUR POLES  
TO BEDDING.



CONTOUR DENSITY

	2-4%
	4-6%
	6-8%
	8-10%

AXIAL PLANE ATTITUDE:  
165/77 E