



OOTSA LAKE GROUP

1a Dacitic Crystal Tuffs; pale green to tan or chalky white. Large K-spar phenocrysts up to 1cm are common and characteristic of this unit. Feldspars are generally fractured. Moderate argillitic alteration-mafics rarely visible.

1b Rhyodacitic Crystal Tuffs; characteristic maroon colour and generally unaltered. Feldspar phenocrysts up to 5mm long are highly abundant. Few quartz phenocrysts. Homogenous - no bedding or welding structures.

1c Welded Rhyodacite Tuffs; commonly shows flow banding and 1-2mm quartz-eyes. Fine-grained pink, grey or yellowish ash matrix.

1d Lapilli Tuffs; light coloured rhyodacitic lapilli fragments with medium to dark grey ash matrix. Minor coal fragments. Dark chalcidony in places.

1e Ash Tuffs; tan to chalky white, slightly to moderately altered. No quartz-eyes. Brecciated and silicified in places.

LEGEND

Brec Brecciated  
 Sil Silification and/or quartz fractures  
 Py Disseminated pyrite  
 Alt Alteration

Soil Sample (Ag ppm, Au ppb)  
 Rock-Chip Sample (Ag ppm, Au ppb)  
 Prefixed by G O

NTS 93/F/3  
 SCALE 1:1000

DATE AUG. 1983 | DRAWN BY RMC/dag | DWG. GC 8021

821708  
 0928/03

WOLF CLAIMS  
 CHOPPER PAD ZONE

GRID I - SOIL, ROCK SAMPLES  
 Ag ppm, Au ppb

Riocanex Inc.

WOLF CLAIMS CHOPPER PAD ZONE

GRID I - SOIL, ROCK SAMPLES  
 Ag ppm, Au ppb

DATE AUG. 1983 | DRAWN BY RMC/dag | DWG. GC 8021