

## RICHTER LITHO STUDY

During the course of mapping on the Richter property in 1989, a total of 141 samples were collected for whole rock analysis. These samples represent all observed lithologies and alteration types seen on the property. The samples were collected during regional mapping of the property, as well as during more detailed mapping programs on the Ridge and Testalinden grids, and in the Ridge Grid ????? trenches. Appendix 1 lists the results of all the litho samples collected. In this listing the samples have been classed into rock and alteration type (in numeric codes), the legend for which is found within the appendix.

Where possible the rock type is taken from field notes. In many cases, however, no identification was made in the field and the rock type has been determined <sup>in conjunction with</sup> by comparison <sup>of the results</sup> with known rock chemistries, ~~combined with~~ <sup>from</sup> descriptions contained in the field notes. Average rock compositions have been determined by plotting histograms for each element for all known unaltered samples of a particular rock type. Samples falling clearly outside the normal distribution for any element have been removed from this data set and reclassified as a different rock type. Some difficulty arises when the data set for a particular rock type is very small, or when there is a wide range in the chemistry of that rock type.

Using the above procedure, the field classification has been modified, in one case adding a previously unrecognized rock type and in another removing a chemically indistinct unit. These changes will be discussed in more detail later in the report. A similar process to that described above was used to assign alteration types to the litho samples.

As a result of the above study, eight different rock types have been recognized on the Richter property. The average composition for each rock type is shown in the following table.

~~Individual rock types will be discussed in detail below.~~

Included in the table are the number of unaltered samples of each rock type collected, since the certainty of the average composition depends on this. Not all compositions have been established to the same degree of certainty. Table

## KOBAU GROUP

### QUARTZITE :

The Kobau Group quartzite is typically a grey to green, fine to medium grained foliated rock. ~~Chemically~~ <sup>It is</sup> difficult to distinguish <sup>between the quartzite and the</sup> granodiorite of the Nelson Plutonic Suite <sup>on the basis of chemistry alone.</sup> The distinction between <sup>however,</sup> unaltered varieties of these rocks should be relatively easy in the field <sup>based on texture.</sup> Subtle chemical variations include a slightly lower percent of Al<sub>2</sub>O<sub>3</sub> and K<sub>2</sub>O, and a slightly higher percent of SiO<sub>2</sub> in the quartzite.

*Only six samples of unaltered quartzite were collected, so the ~~rock type~~ chemical composition is known only approximately.*

### PHYLLITE :

The phyllite unit of the Kobau Group is commonly green in colour (varying shades), fine grained, strongly foliated, and chloritic. The composition of the phyllite varies considerably, as a result of changes in the original rock composition and because of the varying intensity of metamorphism. It may be difficult to identify these rocks based on chemistry alone, because of <sup>the varied chem</sup> this and because of the similarity between these rocks and the gabbroic unit of the Nelson Plutonic Suite. Differences in the percent of Ba and S should allow this distinction. When either the gabbro or phyllite becomes altered significantly, it may be impossible to identify which represents the original host.

*Only three samples of the unaltered phyllite were analysed and the result is very poorly known.*

### MARBLE :

The Kobau Group marble is less widespread than either the quartzite or phyllite. Where seen it is white to pale grey in colour, fine grained, and crystalline. Chemically, these rocks are very distinctive, averaging about 20% in both CaO and SiO<sub>2</sub>. There should be no difficulty identifying these rocks either in the field or by chemical analyses.

## NELSON PLUTONIC SUITE

Three rock types, within the Nelson intrusives, are recognized on the property. Field mapping has suggested that these rock types belong to two separate intrusive events. During the first

episode of intrusion, compositions ranged from dioritic to granodioritic. Field mapping identified a third, fine grained phase (classed as a monzonite), however this study has shown these rocks to be chemically indistinct from the granodiorite. It is believed that the "monzonite" is simply a finer grained version of the same intrusion. While the rocks belong to the same unit, it may be beneficial to identify the finer grained areas, since these may represent the borders <sup>phases or related dykes</sup> of the intrusion. The second intrusive episode is characterized by more mafic intrusives of gabbroic composition.

**GRANODIORITE :**

The granodiorite is typically fine to medium grained, with minor biotite and hornblende crystals visible. Commonly, the rocks are highly weathered. A finer grained, hornblende porphyritic version of the granodiorite is present. These rocks have nearly identical chemical compositions to the coarse grained intrusives and are thought to represent border phases of the intrusion, or possibly related dykes. The average chemical composition of the granodiorite is given in the above table. As discussed above, the composition of the granodiorite and the Kobau Group quartzite is very similar. Distinctions are best made on the percentages of  $Al_2O_3$ ,  $K_2O$  and  $SiO_2$ . When either of the two rocks becomes altered, identification of the original host by chemistry is not possible.

*This composition is based on 18 chemical analyses and believed to be fairly accurate.*

**DIORITE :**

The diorite has been interpreted from field mapping to be a phase of the above intrusion. Typically, these rocks are finer grained and fresher than the granodiorites, and may be hornblende porphyritic. They probably resulted from early fractionation from the same magma. Chemically they are distinct from the granodiorite and also from the later mafic intrusion. Variations between these rock types are most apparent in the percent  $SiO_2$  and  $CaO$ .

**GABBRO:**

*The gabbroic intrusive rocks are typically fine grained, dark green, often with amphibole*

The attached histogram of CaO clearly shows the three rock types. The average percent of CaO of each of the intrusions is shown on the histogram, and it can be seen that they correlate very well with the three populations of data. On a plot of SiO<sub>2</sub>, this distinction is not as clear, although the three intrusives do have markedly different silica content. The lack of distinction could be partly due to the fact that there is a continual transition from granodiorite to diorite, but is probably largely due to the small data set and the biasing towards the granodiorite and gabbroic compositions due to a larger number of samples.