read!

Jolly 82E/3E
825493

Posted: Tue Jul 11, 1989 8:47 AM PDT Msg: HVQA-6011-1000
From: C.F.C.
TO: c.f.c.
Subbject: jolly summary logs

JOL-01 SUMMARY LOG

AZIMUTH 270 deg DIP -45 deg

COLLAR LINE 6 + 30 N
STN 6 + 75 E

0 - 12.6 m.	Overburden
12.6 - 24.1 m	Mafic volcanics
24.1 - 25.6 m	Intermixed volcanics and sediments
25.6 - 28.6 m	Mafic volcanics
28.6 - 35.3 m	Chert
35.3 - 38.1 m	Tuffaceous sediments
38.1 - 38.8 m	Chert
38.8 - 40.2 m	Fault zone (40% chert clasts in clay gouge with local silica flooding; strong graphite bands)
40.2 - 41.7 m	Chert
41.7 - 43.5 m	Intermediate dyke
43.5 - 58.8 m	Chert (may be strongly bx with narrow graphite shears)
58.8 - 73.5 m	Fault zone (Victoria Fault) (very strong graphite shear zone in brecciated chert)
73.5 - 74.4 m	Chert
74.4 - 78.1 m	Fault zone (chert bx and gouge zones, local graphite bands)
78.1 - 80.2 m	Mafic volcanics
80..2 - 81.4 m	Intermediate dyke
81.4 - 85.0 m	Diorite
85.0 - 86.0 m	Fault zone (strongly silic'd diorite, bx)
86.0 - 105.0 m	Diorite
105.0 - 106.8 m	Intermediate dyke
106.8 - 150.0 m	Diorite
150.0 m	E. O. H.

Tested the Victoria shear zone to the north of the Victoria #2 adit. Found a strong graphitic shear hosted in cherts.

JOL-02 SUMMARY LOG

AZIMUTH 270 deg DIP -45 deg

COLLAR LINE 5 + 10 N
STN 6 + 85 E

0 - 9.4 m	Overburden
9.4 - 12.9 m	Mafic volcanics
12.9 - 13.5 mm	Fault zone (green clay gouuge with 50% silic. bx clasts)
13.5 - 14.9 m	Mafic volcanics
14.9 - 18.3 m	Faault zone (strongly bx, silicified, 2-5% py)
18.3 - 42.8 m	Mafic volcanics
42.8 - 46.9 m	Tuffaceous sediment
46.9 - 47.6 m	Limestone
47.6 - 49.4 m	Tuffaceous sediment
499.4 - 55.5 m	Chert/Fault zone (Victoria fault ?) (strongly bx chert clasts in black, graphitic, py-rich gouge)
55.5 - 70.2 m	Mafic volcanics
70.2 - 71.1 m	Diorite dyke
71.1 - 80.4 mm	Mafic volcanic
80.4 - 80.8 m	Cherty argillite
80.8 - 81.8 m	Mafic volcanic
81.8 - 883.1 m	Fault gouge (white clay gouge with 5% qtz frags)
83.1 - 105.8 m	Diorite

105.8 m	E. O. H.

Tested the Victoria shear zone in the area between the Victoria #1 and #2 workings. Intersected several major faults. The graphitic brecciated fault zone hosted in cherts at 49.4 - 55.5 m is believed to be the Victoria shear zone.

JOL-0003 SUMMARY LOG

AZIMUTH 270 deg DIP -45 deg

COLLAR LINE 4+00 N
STN 6+75 E

0 - 6.8 m	OVERBURDEN
6.8 - 8.3 m	CHERT/FAULT ZONE Strongly bx with local gouge zones.
8.3 - 9.5 m	TUFFACEOUS SEDIMENTTS
9.5 - 11.8 m	MAFIC VOLCANICS
11.8 - 12.7 m	CHERT
12.7 - 113.4 m	TUFFACEOUS SEDIMENTS
13.4 - 18.4 m	MAFIC VOLCANICS
18.4 - 20.0 m	INTERBEDDED VOLCANICS AND SEDIMENTS
20.0 - 42.0 m	MAFIC VOLCANICS
42.0 - 43.0 m	INTERBEDDED VOLCANICS AND SEDIMENTS
43.0 - 60.2 m	MMAFIC VOLCANICCS
60.2 - 62.4 m	TUFFACEOUS SEDIMENTS
62.4 - 63.2 m	MAFIC VOLCANICS
63.2 - 64.8 m	CHERT
64.8 - 73.3 m	DIORITE DYKE Siliceous, bleached, weak sericite alt'n.
73.3 - 73.9 m	FAULT ZZONE Very strong clay gouge with 20% alt'd diorite clasts.
73.9 - 75.3 m	CHERT/TUFFACEOUS SEDIMENTS
75.3 - 79.8 m	MAFIC VOLCANICS
79.8 - 91.5 m	FAULT ZONE (Vicctoria faault) Intensely bx chert, minor graphite bands and gouge zones.
91.5 - 95.5 m	MAFIC VOLCANICS
95.5 - 101.0 m	FAULT ZONE (Victoria fault) Intense gouge zone.
101.0 - 111.1 m	DIORITE Bleached, siliceous, sericitized.
111.1 - 112.9 m	DIORITE Propylitic alt'n.
112.9 - 113.4 m	DIORITE/FAULT ZONE Bleached, strong sericitization, ssilc'n.
113.4 - 114.1 m	DIORITE Propylitic alt'n.
114.1 - 117.0 m	DIORITE/FAULT ZONE Bleached, strong silic'n.
117.0 - 126.9 m	DIORITE
126.9 - 127.5 m	DIORITE/FAULT ZONE Bleachhed, silicified, minor gouge zones and v strong sericitization.
127.5 - 138.8 m	DIORITE
138.8 - 140.6 m	DIORITE/FAULT ZONE Strong bx, silic'n, 5% py in bx matrix.
140.6 - 150.3 m	DIORITE
150.3 - 150.7 m	GRANODIORITE DYKE
150.7 - 165.2 m	DIORITE
165.2 m	E. O. H.

Tested the Victoria Fault near the Victoria #1 adit. Intersected this faultt at 79.8 - 91.5 m as well as intersecting a major steep east dipping fault zone with assoociated silicified diorite, trending up the main creek and exposed in the Victoria #1 adit.

JOLLY SUMMARY LOG: HOLE 04

AZIMUTH 270 deg DIP -45 deg

COLLAR LIINE 3+00 N
STN 2+12 E

0- 12.2 m	OVERBURDEN
12.2 - 20.3 m	MAFIC VOLCANICS
20.3 - 48.4 m	DIORITE
48.4 - 51.3 m	FAULT ZONE Siliceous, minor green mica, locally strong bx and gouge zones.
51.3 - 54.8 m	HORNFELS-ARGILLITE
54.8 - 55.8 m	FAULT ZONE Strongly bx, sericite alt'n and gouge zones.
55.8 - 59.7 m	ARGILLITTE
59.7 - 60.8 m	FAULT ZONE Strongly bx, strong sericite alt'n.
60.8 - 62.8 m	INTERBEDDED VVOLCS/ARGILLITE
62.8 - 64.4 m	FAULT ZONE Strong talc alt'n.
64.4 - 65.2 m	MAFIC VOLCANICS
65.2 - 66.1 m	FAULT ZONE Graphitic-siliceous, banded.
66.1 - 66.8 m	MAFIC VOLCANICS
66.8 - 87.8 m	FAULT ZONE Graphitic-siliceous, banded, bx.
87.8 - 92.2 m	FELDSPAR PORPHYRY DYKE Siliceous
92.2 - 99.0 m	FAULT ZONE Graphitic-siliceous, banded, bx.
99.0 - 105.0 m	FELDSPAR PORPHYRY DYKE Siiliceous
105.0 - 135.0 m	FAULT ZONE Graphitic-siliceous, banded, bx.
135.0 - 138.6 m	CHERT
138.6 - 140.6 m	FAULT ZONE Graphitic-siliceous, banded, bx.
140.6 - 1446.3 m	CHERT
146.3 - 148.5 m	MAFIC V

JOLLY SUMMARY LOG: HOLE 05

AZIMUTH 270 deg DIP -45 deg

COLLAR LINE 5+00 N
STN 3++17 E

0 - 27.4 m	OVERBURDEN
27.4 - 31.2 m	DIORITE Silicified.
31.2 - 44.1 m	FAULT ZONE Strong talc alt'n, locally siliceous, bx.
44.1 - 477.7 m	DIORITE
47.7 - 51.8 m	FAULT ZONE Graphitic, bx cherty argillite.
51.8 - 68.6 m	FAULT ZONE Stronng talc alt'n, locally siliceous, bx.
68.6 m	E. O. H.

Tested the north trending fault contact of the diorite and ssediments to the north of the Gold Standard workings. The hole was lost in the strong talc fault zone at 68.6 m.

Posted: Tue Jul 25, 1989 9:16 AM PDT Msg: PVQA-2333-1000
From: C.F.C.
TO: c.f.c.
Subject: month end

JOL-06 SUMMARY LOG

AZIMUTH 000 deg DIP -45 deg

COLLAR LINE 9+50 N
STN 7+25 E
ELEV 1094 m

0 - 12.2 m	OVERBURDEN
12.2 - 20.0 m	DIORITE
20.0 - 32.7 m	ALT'D DIORITE (Intense sericitization + py + local silic'n.)
32.7 - 33.6 m	FAULT ZONE (Intense clay-sericite gouge, 10% diss py.)
33..6 - 34.4 m	FINE GRAINED INTERMEDIATE DYKE
34.4 - 47.9 m	ALT'D DIORITE (Intense serpentinite, mod silic'n, sericitiz'n. Local gougee and bx zones, 5-10% py.)
47.9 - 52.4 m	ALT'D DIORITE (Mod serpentine, minor sericitiz'n.)
52.4 - 60.8 m	ALT'D DIORITE (V. strong sericitiz'n/talc, minor serp, mod-sstrongly bx, local gouge zones.)
60.8 - 69.8 m	DIORITE
69.8 - 71.4 m	FAULT ZONE (Strongly alt'd bx diorite and fault gougge.)
71.4 - 98.9 m	ALT'D DIORITE (Strong talc-sericite alt'n, locally silicified or serpentinized. May be mod-strongly bx.)
98.9 - 104.5 m	FELDSPAR PORPHYRY DYKE
104.5 - 111.9 m	DIORITE
111.9 mm	E. O. H.

Tested magnetic low under the placer channel. Intersected over 100 metres of very strong alteration (serp, seric, silic) before hole was lost due to caving.

Posted: Tue Jul 25, 1989 9:20 AM PDT Msg: XVQA-3333-1000
From: C.F.C.
TO: c.f.c.
Subject: jolly summ logs

summary logs

JOL-07 SUMMAARY LOG

AZIMUTH 180 deg DIP -45 deg

COLLAR LINE 10+50 N
STN 3+00 E
ELEV 1125 m

0 - 94.4 m	OVERBURDEN
94.4 - 111.3 m	MAFIC VOLCANICS (carb, chl alt'n, locally siilic)
111.3 - 113.7 m	IMPURE LIMESTONE
113.7 - 114.9 m	MAFIC VOLCANICS (chl-actinolite alt'n)
114.9 - 118.5 m	MAFIC VOLCANICS (intense silic'n)
118.5 - 120.0 m	FAULT ZONE (graphitic-siliceous, banded, 1% py)
120.0 - 121.9 m	MAFIC VOLCANICS (strong ssilic'n)
121.9 - 125.6 m	FAULT ZONE (graphitic-siliceous, banded, 2% py)
125.6 - 130.7 m	MAFIC VOLCANICS
130.7 - 136.6 m	IMPURE LIMESTONE
136.6 - 166.2 m	MAFIC VOLCANIC (chl-carb aalt'n)
166.2 - 177.2 m	FAULT ZONE (graphitic-siliceous, banded, bx, 2% py)
177.2 - 177.5 m	INTERMEDIATE DYKE (hematitic)
177.5 - 183.4 m	FAULT ZONE (graphitic-siliceous, banded, bx, 2% py)
183.4 - 184.4 m	INTERMEDIATE DYKE (strong siliic'n, 2% py)
184.4 - 186.8 m	FAULT ZONE (graphitic-siliceous, banded, bx, 2% py)

186.8 - 187.8 m	INTERMEDIATE DYKE (strong siliic'n, hematitic, 2% py)
187.8 - 193.8 m	FAULT ZONE (graphitic-siliceous, banded, bx, 1% py)
193.8 - 196.7 m	INTERMEDIATE DYKE (siliceous, weak chl alt'n, 2% py)
196.7 - 197.6 m	CHERT
197.6 - 210.3 m	FAULT ZONE (graphitic-siliceous, banded, bx, 2% py)
210.3 - 210.9 m	IMPURE LIMESTONE
210.9 - 213.0 m	FAULT ZONE (graphitic-siliceous, banded, bx, 2% py)
213.0 - 215.2 m	IMPURE LIMESTONE
215.2 m	E. O. H.

Tested the mag low to the west of JOL-06, drilling from the north. The drill hole remained in sediments & volcanics throughout its length, intersecting a major vertical graphitic shear zone, on trend with the serpentinite zone in JOOL-06.

Posted: Tue Jul 25, 1989 9:22 AM PDT Msg: GVQA-4333-11000
From: C.F.C.
TO: c.f.c.
Subject: jolly summ log

JOL-08 SUMMARY LOG

AZIMUTH 180 deg DIP -45 deg

COLLAR LINE 11+50 NN
STN 5+35 E
ELEV 1116 m

0 - 97.5 m	OVERBURDEN
97.5 - 103.4 m	INTERBEDDED MAFIC VOLCANICS/CHERTY SEDS
103.4 - 104.3 m	FAULT ZONE/MAFIC VOLCANICS (silicified, bx, narrow gdior dyke in core)
104.3 - 107.0 m	INTERBEDDED MAFIC VOLCANICS/CHERTY SEDS
107.0 - 108.0 m	CHERT
1008.0 - 110.0 m	MAFIC VOLCANICS
110.0 - 110.9 m	FELDSPAR PORPHYRY DYKE (intensely siliceous, 2-5% py)
110.9 - 112.8 m	CHERT
112.8 - 115.222 m	MAFIC VOLCANICS
115.2 - 118.0 m	FAULT ZONE/FELDSPAR PPORPHYRY DYKE (intense silic'n, bx, narrow gouge zones)
118.0 - 119.0 m	MAFIC VOLCANICS
119.0 - 1255.4 m	DIODRITE (or med gr mafic volc?)
125.4 - 131.8 m	CHERT & CHERTY TUFFS
131.8 - 136.8 m	DIORITE/FAULT ZONE (strong silic'n, sericitization)
136.8 - 139.5 m	FAULT ZONE (fault gouge with bx frags of seric diorite)
139.5 - 141.6 m	DIORITE/FAULT ZONE (strong seric, serp alt'n, 5-10% qtz veinlets)
141.6 - 148.7 m	DIORITE/FAAULT ZONE (strong seric, talc alt'n, minor silic'n)
148.7 m	E. O. H.

Tested the mag low between JOL-06 & JOL-07, drilling from the north. The hole was lost, due to squeezing, shortly after intersecting a vertical, strongly alttered (serp/seric) zone in the diorite.

Posted: Tue Jul 25, 1989 9:23 AM PDT Msg: UVQA-4333-1000
From: C.FF.C.
TO: c.f.c.
Subject: jolly suumm log

JOL-09 SUMMARY LOG

AZIMUTH 360 deg DIP 45 deg

COLLAR LINE 8+00 N
STN 5+17 E
ELEV 1115 m

0 - 36.6 m	OVERBURDEN

36.6 - 115.4 m	MAFIC VOLCANICS (strong carb alt'n)
115.4 - 130.5 m	DIORITE (mod chl alt'n)
130.5 - 150.8 m	FAULT ZONE/ALTERED DIORITE (strong talc-sericite alt'n, bx, local gouge zones))
150.8 - 152.4 m	DIORITE DYKE (mod chl alt'n)
152.4 - 160.0 m	FAULT ZONE/ALTERED DIORITE (strong sericite-talc alt'n, bx, locally siliceouus, minor green mica)
160.0 - 163.4 m	FELDSPAR PORPHYRY DYKE (ssilicified, mod chl, seric alt'n, minor py)
163.4 - 174.0 m	FAULT ZONE/ALTERED DIORITE (strong sericite alt'n, locally serpentinized or silicified , bx)
174.0 - 175.2 m	FELDSPARR PORPHYRY DYKE (strong chl alt'n)
175.2 - 178.9 m	FAULT ZONE/ALTERED DIORITE (strong sericite alt'n & fault gouge)
178.9 m	E. O. H.

The hole was drilled from the south in order to test the mag low along section with JOL-08, which was lost before entering the core of tthe zone. This hole was also lost, about 40 m into the zone, before the core of alteration was reached.

Posted: Tue Jul 25, 1989 9:34 AM PDT Msg: QVQA-8333-1000
From: C.F.CC.
TO: c.f.c.
Subject: brrenda month end

JULLY MONTH END

JOLLY - 628

- 9 NQ diamond drill holes (a total of 1299 metres) were drillled in July on three different targets
- JOL-01, 02 & 03 tested the Victoria fault zone. Although the structure is a significant width, gold values are generally low. Anomaloous gold values (to 2290 ppb over 1 m) do occur w/narrow quartz veins.
- JOL-04 & 05 were drilled to test the western fault boudary of the diorite and sediments. Hole 4 intersected a 92 metre wide graphitic fault zone with only slightly elevated gold and silver values (to a maximum of 71 ppb and 1.2 ppm, respectively). Hole 5 was lost in a strong talc fault zone at a depth of 68 metres. Again, gold and silver values were only slightly elevated (to a maximum of 97 ppb and 1.4 ppm, respectively).
- JOL-06, 07, 08 & 09 were drilled to test the E-W trending magnetic low under the pacer channel. Drilling intersected a major fault zone with intense alteration (sericite, talc, serpentinite). Because of difficulties in drilling, the entire fault zone was nevver intersected. No results have been received from any of these holes.
- Mapping & rock sampling on the property confirmed that, to date,,, all known gold mineralization occurs in narrow quartz veins. Values to 221 g/tonne were obtained from a grab sample off a narrow quartz vein at the Old England workings.
- 10 line kilometres of geophysics (magnetometry, VLF-EM) was completed on the extension of the grid. No signifacant targets resulted from this program, although thee existing E-W mag low was confirmed and expanded by the program.

WART - 643

The property was located and briiefly exxamined in July. Access to the property is much better from Merritt so work will be carrried out at the same time as work on Clapper.

RICHTER - 656

Reconnaissance mapping was completed on the property and three potential targets have been identified. The first target is in the form of a 2200 ppb Au returned from a silicified quartzite adjacent to intrusive and gossan. The second target has not, as yet, returned anomalous values but is an interesting melange of silicified rocks & quartz veins with the possibilty of ultramafic being included in the package. The most interesting target is an area with 2 intersecting quartz veins with up tto 6m widths. The veins run along granodiorite intrusives. One 350m length of intrusive is highly altered and gossanous and has returned a 6800 ppb Au value (0.232 oz/ton assay). A tight grid is, at present, being established over the area and detailed mapping and sampling are underway.

BRENDA RECCE - 6244, 658

Heavy mineral sampliing in the Kelowna area has returned several anomalous drainages in the Westbank region. These wiill be followed up and compiled.

Several properties were examined during the month. Tertiary systems include the Golden Plug which is a larrge felsic intrusive centre with extensive clay alteration and anomalous AAs, Zn values, the Astro pproperty (owned by Minequest) was visited and the anomalous Au bearing shear zone was examined. Also Geotec's property on the old southern limit of the Astro claims are being investigated.

The Dividend property was visited and numerous As, Po, Cpy skarns were sampleed with dissappointing results (max 0.4% Cu, 4.3 ppm Ag and 112 ppb Au). Meanwhile, the Lakeview-Dividend skarn visited earlier in the season returned values up to 127 g/t Au (3.70 oz/t).

Several quartz veins in the Richter area were visited including Mac Siccar, Stemwinder, Powis and Tinhorn. Values up to 123 g/t Au and 83 g/t AAg have been returned from these veins which are similar to those being explored on Testalinden Creek. An interesting sample taken 10m above the Stemwinder workings returned a Au value of 2 ppb which may reflect surface leaching.

```
Command?  scan

  No new messages.

Command?
```