

Foremore Project; Geology

LEGEND

LITHOLOGIES

Intrusive Rocks:

- More Creek Pluton, basic gneiss and local nepheline, mica phases and zirconiferous
- Unmetamorphosed Tertiary Eryonite, Monzonite, Gneiss and Siltstone
- Dorono-Mississippi(?) Diorite / Gabbro

Stratified Rocks:

Upper Tertiary:

- Unmetamorphosed Group Volcanics and Volcanic Conglomerates

Metamorphic Rocks:

- Felsic Volcanics
- Mafic and Intermediate Volcanics and Flow
- Limestone and associated Limestone with minor Chert, Argillite and Turf
- Green and Mason Turf and Flow Breccias
- Carbonaceous Argillite, Cherty Argillite and Chert Pieces Conglomerates

Diverse Metasiltstone:

- Crinoid Limestone
- Quartz and/or Barite Gneiss
- Nepheline and Chlorite Phyllite and Chlorite Siltstone
- Felsic Turf
- Intermediate to Mafic Volcanics
- Argillaceous Phyllite and Black Siliceous Phyllite
- Silver or White Phyllite and Quartziferous Siltstone

Geochronology


- Rock sample
- Soil sample
- Drill sample

Geophysics

- Strong UTEM conductor
- Moderate UTEM conductor
- Weak UTEM conductor

Other Symbols:

- Drill hole



North Boulder Field (NBF)
+ 100 VHMS boulders
 Layered Avg (29) = 1.1g/t Au, 87g/t Ag, 10.2% Zn, 3.5% Pb
 Feeder Avg (12) = 1.5g/t Au, 167g/t Ag, 6.2% Zn, 2.3% Cu

Discovery Boulder
 5.2 oz/T Au (Cominco 1987)

North Zone Targets

Side Glacier BF (SGBF)
 Avg (8) = 8.0g/t Au, 7.6% Zn, 1.7% Pb

East Boulder Field (EBF)

See schematic cross section


Westmore Au Targets

Hanging Valley Zones
 2.73 oz/T Au, 1.0 oz/T Ag, 1.1% Cu
 11.7% Pb, 0.8 Zn for 1.0m
 (Cominco 1989)

See schematic cross section

South Boulder Field (SBF)
 Avg (53) = 102g/t Ag, 9.4% Zn, 1.7% Pb

Nunatak Zone



Roca Mines Inc.

**Foremore Property
 Compilation Map**

Date	Aug 27, 2002	Scale	1:20,000	Figure
UTM Zone	9 - NAD83	Mining District	LIARD	
N.T.S.	104G/2, 104B/14, 15	State/Province	BC	

883009