SMGATIJAKIOA./VAP

1710 - 609 GRANVILLE ST PO BOX 10363 VANCOUVER BC CANADA V7Y 1G5 (604) 683-7265 FAX 683-5306 BBS 683-7206

NO. 210 (1994) **NOVEMBER 2, 1994**

George Cross News AEEEVE Debushed Daily ALL REPRODUCTION

"Reliable Reporting" WESTERN CANADIAN INVESTMENTS

SUBSCRIPTION RATE 3 1994 \$315.00 PER YEAR

COPYRIGHT

E. M. P. R.

NO. 210 (1994) **NOVEMBER 2, 1994**

BOOKER GOLD EXPLORATIONS LTD. [BGE-V] 4,357,035 SHS.

HEARNE HILL PROJECT UPDATE - Malcolm Macdonald, director, **Booker** Gold Explorations

Ltd., report further developments at the Hearne Hill project 65 km. northeast of Smithers and 15 km north of the Bell Copper Mine in the Babine Lake District of B.C. The property is 100% owned by Booker Gold, subject to a 4% NSR which can be bought out.

The company has received drill log information from DDH 94-5. The hole is mineralized on fractures and contains disseminated mineralization throughout. Mineralization is primarily the copper sulphide chalcopyrite and some bornite, with limited molybdenite on fractures. Secondary biotite and potassium feldspar alteration is extensive. Of particular interest is the repeated intersections of breccia from a depth of 100 to 216 metres. The mineralization in the breccia zones is commonly chalcopyrite with some pyrite. The breccia zones are mineralized in fractures and open space filling between volcanic and occasional biotite feldspar porphyry fragments. The chalcopyrite is often found as globs within open spaces. The brecciated zones are strongly siliceous and at 216 metres quartz sericite molybdenite and bornite were noted with chalcopyrite in open space filling and fractures. At the bottom of the hole there is a sharp contact between rhyodacite and biotite feldspar porphyry. The rhyodacite is siliceous and biotite altered with veinlets of chalcopyrite. The biotite feldspar porphyry is extremely well mineralized with chalcopyrite and is sericite and clay altered. (SEE GCNL NO.207, Oct.28/94, P.2 FOR PREVIOUS PROJECT INFORMATION)

M 6