

10:35 Red Mountain (Clone), BC*Ed Kruckowski, Teuton Resources Corporation*

The Clone property, owned by Teuton Resources Corp. and Minvita Enterprises Ltd. is located 16 km southeast of Stewart, British Columbia in the Skeena Mining Division. The property covers an area of Hazelton pyroclastic volcanic rocks in contact with a variety of intrusive plutons associated with the main Coast Range Batholith.

During 1995, the property was explored by 513.8 m of trenching, ground geophysics and 1070 m of BTW drilling in 13 holes. During 1996, the property was explored by 1312.8 m of trenching in 141 trenches, ground geophysics and 11,487.1 m of both BTW and NQ2 drilling in 113 holes.

Mineralization within the main area of interest consists of two different and distinct types. The mineralization is hosted by steeply dipping sub-parallel, en echelon, shear controlled veins and stockwork with a northwesterly trend. The first type of mineralization is dominated by pyrite plus/minus arsenopyrite plus/minus chalcopyrite with gold and cobalt values within chlorite, schistose lapilli tuffs and the second by massive hematite veins with associated chlorite and calcite-quartz stockworks within broad zones of hematite-chlorite-K-feldspar altered rocks. Specularite, chalcopyrite, magnetite and locally visible gold are associated with the hematite-dominated mineralization. The sulfide-dominated mineralization prevails along strike lengths up to 2 km. The hematite-dominated structures have less defined walls but show good strike lengths as well.

There is some evidence to suggest that the sulfide mineralization is later and some overprinting along the hematite bearing structures has occurred. Based on the gold-arsenic-cobalt geochemical association, a potential strike length of at least 5.5 km is indicated for the main gold bearing structures.

10:50 Hearne Hill, BC*Gordon Weary, Booker Gold Explorations Ltd.*

Booker Gold Explorations Ltd. is currently involved in an aggressive drill program on the Hearne Hill property, located in west-central British Columbia. Drilling during the winter of 1997 will determine the size and extent of enriched copper-gold breccia zones discovered in 1996 and attempt to locate new areas of high-grade mineralization.

The Hearne Hill property is located 65 km northeast of Smithers, British Columbia. Hearne Hill is a classic porphyry copper-gold deposit associated with rocks of the Babine Igneous Suite. The presence of chalcopyrite-cemented breccia bodies at Hearne Hill is unique among the Babine deposits. Former open pit producers in the area are the Bell Copper Mine, 15 km to the south (128 m tonnes of 0.48% Cu, 0.30 g/t Au) and the Granisle Mine, 18 km to the south (94 m tonnes 0.43% Cu, 0.12 g/t