

CIM Dist 6 Kamloops Sept 6-8/90

CIM Bull V83 July 90

675240

Tulsequah
Chief
104K/12

Coffee Break — 10:00 a.m.

Paper No. 23 — 10:30 a.m.

Cominco-Redfern Tulsequah Chief Massive Sulphide Deposit — Northwest British Columbia.

R. AULIS, Cominco Ltd., Vancouver, British Columbia

The Tulsequah Chief property is located near the confluence of the Tulsequah and Taku Rivers in the Coast Range Mountains of British Columbia, 100 km south of Atlin, British Columbia, and 65 km northeast of Juneau, Alaska. The property was first staked in 1923 following the discovery of a high-grade lens of pyrite, barite, sphalerite, galena and chalcopyrite. Cominco Ltd. acquired the deposit in 1946 and placed it and the adjacent Big Bull deposit into production in 1951 at a rate of 530 tons/day. The mine was closed in 1957 due to low metal prices.

Production from 1951-1957 was 625 781 tons from the Chief and 403 308 tons from

46

the Bull at a combined average grade of 0.11 oz/t Au, 3.69 oz/t Ag, 1.59% Cu, 1.54% Pb and 7.0% Zn. At shutdown, ore reserves in the Chief to the 5100 level were estimated at 780 000 tons at 0.07 oz/t Au, 2.9 oz/t Ag, 1.3% Cu, 1.6% Pb and 7.0% Zn. Recent drilling has discovered two new sulphide lenses and extended others. Redfern has calculated a potential geological reserve, including previous reserves, to the 3500 level of 5.8 million tons at 0.08 oz/t Au, 2.9 oz/t Ag, 1.6% Cu, 1.3% Pb, 7.0% Zn. Average spacing of drill hole penetrations is too widespread to infer hole-to-hole correlation with a high degree of confidence.

The Tulsequah Chief deposit occurs in a northeasterly-striking, west-dipping sequence of Pre-Permian, submarine, volcanic-sedimentary rocks located on the west limb of a north-plunging anticline. The anticline is delineated by a mixed limestone, chert, clastic sequence containing Pennsylvanian-Permian fossils. This sequence occurs stratigraphically above the Chief deposit. Pre-Permian rocks consist primarily of basalt-andesite volcanics with lesser dacite-rhyolite, pyroclastics, clastics, limestone and chert. All rocks are intruded by Paleozoic diorite and dacite, and Tertiary rhyolite plugs, stills and dykes. A major regional fault cuts off the west extension of the Tulsequah Chief stratigraphy.

The Tulsequah Chief deposit is located along the base of a thick lenticular mass of dacite-rhyolite pyroclastics near the contact with an underlying thick sequence of basalt-andesite pyroclastics and flows. The deposit is broken into four blocks by north-south striking, steeply dipping faults, all of which were initially synvolcanic growth faults which created a seafloor graven that was active during deposition of the sulphide lenses.

Mineralization occurs in conformable lenses consisting of pyrite (15% to 80%) with varying amounts of sphalerite, galena, chalcopyrite, gold, silver, barite and gypsum. These lenses occur along a 700 m strike length in a lithologic package known as the mineral horizon. The mineral horizon consists of altered, intercalated dacite-rhyolite tuffs, muds, cherty tuffites, and cherts intermixed with altered dacite-rhyolite lapilli tuffs.

Alteration in the mineral horizon consists primarily of sericite and pyrite (10% to 80%). Weak sericite-pyrite (<5%) alteration and patchy anastomosing zones of silica veins and pervasive silicification extend for distances up to 30 m above the mineral horizon into the overlying dacite-rhyolite pyroclastic package, indicating hydrothermal activity continued after the main phase of sulphide deposition. The mineral horizon is underlain by a discordant alteration pipe which is centred on the graben, crosscuts stratigraphy, can be traced on surface for 1 km and occurs primarily in andesite volcanics. Alteration in the pipe is concentrically zoned outward from a sericite-pyrite (15% to 25%) core through a transition zone of biotite (15% to 25%) to an outer zone of chlorite-pyrite (5% to 10%). An adjacent, separate alteration pipe, 400 m to the west, formed slightly later at a higher stratigraphic level. No mineralization of significance has been found to date associated with this second pipe.

The intimate spatial relationship of the mineralized lenses with the same package of volcanic rocks indicates that sulphide formation was an integral part of, and related to volcanism. The Tulsequah Chief is typical of the rhyolite associated massive sulphide class of deposits with its closest analogue, Westmin's HW deposit on Vancouver Island.

new zone described
verbally at talk
~150ft of MS
cut in DDH below
main zone.